

Indiana Collegiate Action Network

Results of the
**Indiana College
Substance Use Survey
2012**



Results of the Indiana College Substance Use Survey 2012

by

Rosemary King, M.P.H.
Mi Kyung Jun, Ph.D.

Indiana Prevention Resource Center
501 N. Morton St., Suite 110
Bloomington, Indiana 47404
www.drugs.indiana.edu

Online survey and report cover designed by Junghun Lee, M.S.

Funded by the Indiana Family and Social Services Administration,
Division of Mental Health and Addiction
through a contract with the Indiana Collegiate Action Network

Recommended citation

King, R.A., & Jun, M.K. (2012). *Results of the Indiana College Substance Use Survey – 2012*.
Bloomington, IN: Indiana Prevention Resource Center, Indiana University.

TABLE OF CONTENTS

List of Tables	ii
List of Figures	iii
Introduction	1
Methodology	3
Key Findings	6
Prevalence and Frequency of Substance Use	6
Consequences of Substance Use	18
Risk Factors	21
Gambling Behaviors	28
Characteristics of Survey Participants	32
References	34
Frequency Tables	37

LIST OF TABLES

Table 1. Select Demographic Characteristics of Survey Respondents, 2009 - 2012	2
Table 2. Response Rate, 2012	4
Table 3. Number of Usable Questionnaires, 2012	5
Table 4. Percentage of Students Reporting Use of Select Substances in the Past Year, by Select Demographic Characteristics, 2012.....	8
Table 5. Percentage of Students Reporting Use of Select Substances in the Past Month, by Select Demographic Characteristics, 2012.....	9
Table 6. Percentage of Students Reporting Binge Drinking in the Past Month, by Select Demographic Characteristics, 2012	11
Table 7. Percentage of Students Reporting Binge Drinking in the Past Two Weeks, by Select Demographic Characteristics, 2012	11
Table 8. Percentage of Students Reporting Prescription Drug Misuse in the Past Six Months, by Select Demographic Characteristics, 2012.....	13
Table 9. Percentage of Students Who Drank Alcohol in Past Six Months, Who Indicated Select Reasons for Drinking Alcohol, 2012 (n = 5,651)	17
Table 10. Percentage of Students Who Drank Alcohol in the Past Six Months, Who Reported Select Consequences as a Result of Their Drinking, 2012 (n = 5,651).....	19
Table 11. Percentage of Students Who Used Drugs in the Past Six Months, Who Reported Select Consequences as a Result of Their Drug Use, 2012 (n = 1,523)	20
Table 12. Average Age of First Time Use of Select Substances, 2012.....	22
Table 13. Three Easiest Ways of Getting Alcohol as Reported by Students Who Are Under 21 and Who Drink Alcohol, 2012 (n = 2,827).....	24
Table 14. Mean Number of Drinks Respondents Thought the Typical Student Consumed the Last Time He/She “Partied/Socialized”, 2012 (range = 0 – 30).....	27
Table 15. Percentage of Students Who Thought Other Students on Campus Would Approve or Disapprove of Select Behaviors, 2012	27
Table 16. Percentage of Students Reporting Gambling Behaviors in the Past Year, 2012	29
Table 17. Percentage of Students Who Gambled in the Past Year, Who Reported Select Consequences as a Result of Their Gambling, 2012 (n = 2,811)	31
Table 18. Demographic Characteristics of Survey Participants, 2012	32

LIST OF FIGURES

Figure 1. Frequency of Binge Drinking in the Past Two Weeks, 2012	11
Figure 2. Percentage of Students Who Drank Alcohol in Past Six Months, Who Combined Alcohol with Energy Drinks or Other Stimulants, 2012 (n = 5,651)	14
Figure 3. Percentage of Students Who Drank Alcohol in Past Six Months, Who Reported Select Categories as the Usual Type of Alcohol They Drank in the Past Month, 2012 (n =5,651).....	15
Figure 4. Percentage of Students Under 21 Who Drink Alcohol, Who Reported Purchasing Alcohol from a Retailer in the Past Six Months, 2012 (n = 2,827).....	24
Figure 5. Percentage of Students Under 21 Who Have Purchased Alcohol from a Retailer, Who Reported Select Methods as the Most Common for Purchasing Alcohol, 2012 (n = 287).....	25
Figure 6. Percentage of Students Under 21 Who Reported Purchasing Alcohol from Retailers, Easiest Place to Purchase Alcohol, 2012 (n = 287)	25
Figure 7. Percentage of Students Who Reported Weekly or More Frequent Gambling in the Past Year, by Gambling Category, 2012.....	30
Figure 8. Most Frequent Gambling Location (Other Than Online or Lottery) in Past Year, 2012.....	30

INTRODUCTION

Substance use by college students has long been recognized as a serious problem on our nation's campuses, with alcohol being the most widely used substance. In 1998, the National Institute on Alcohol Abuse and Alcoholism undertook a comprehensive assessment of drinking on college campuses, concluding that social and environmental influences present on most campuses have created a culture of drinking, where alcohol consumption has become a rite of passage. Drinking to excess (binge drinking) is widespread among college students, and has serious consequences (NIAAA, 2002). The rates of alcohol and drug use have remained relatively steady over the past decade, with approximately two-thirds of college students nationwide using alcohol in the past month, approximately two-fifths engaging in binge drinking in the past two weeks, and approximately one-fifth using illicit drugs in the past month (Johnston, O'Malley, Bachman, & Schulenberg, 2012).

In 2006, the Indiana Collegiate Action Network (ICAN) provided funding for the first uniform assessment of substance use across institutions of higher education in Indiana, utilizing the Core Alcohol and Drug Survey. Beginning in 2008, the Indiana Division of Mental Health and Addiction (DMHA) has awarded funding to ICAN to continue the statewide collection and analysis of data on the prevalence of substance use by Indiana college students. ICAN, in conjunction with the Indiana Prevention Resource Center (IPRC) developed a new survey instrument in 2009, called the Indiana College Substance Use Survey, to collect information on items of interest to Indiana colleges.

This report summarizes the findings of the fourth annual Indiana College Substance Use Survey that was conducted by the Indiana Prevention Resource Center in the spring of 2012. Findings on the prevalence of alcohol, tobacco and other drug use, consequences of use, alcohol availability, and student perceptions of peer behaviors are presented. The findings have been analyzed across gender and age variables. National prevalence data from the Monitoring the Future Study have been included to facilitate comparison between the participating Indiana college students and the national rates, where available. The Monitoring the Future data are the most recent national data (2011) on college students' prevalence rates of substance use available at the time of this report.

Results from previous years are not presented in this report because the participating schools vary from year to year. The small number of schools participating in the survey and the absence of a random sampling technique make it unsuitable to generalize the findings to Indiana college students overall. Therefore, the data should not be compared to identify any trends in behaviors over time. The following table provides some demographic data of the survey participants each year. The previous years' reports can be found at www.drugs.indiana.edu/data-survey_icsus.html.

Table 1. Select Demographic Characteristics of Survey Respondents, 2009 - 2012
(Values are percentages)

	2009	2010	2011	2012
Gender				
Male	42.9	36.8	36.2	36.9
Female	57.1	63.2	63.8	63.1
Age				
Under 21	53.4	46.1	46.4	53.9
21 or older	46.6	53.8	53.6	46.1
Race				
White	86.9	87.9	86.4	84.4
Black/African American	2.3	3.7	3.9	3.4
Asian	3.7	2.6	4.3	7.2
Other	6.4	5.2	4.8	4.3
Ethnicity				
Hispanic	5.5	3.2	3.1	3.7
Student status				
Full-time	95.6	89.5	92.0	97.0
Location of residence				
On-campus	61.9	50.3	45.1	54.8
# of usable surveys	4,885	5,928	5,304	6,968

METHODOLOGY

The Indiana College Substance Use Survey was developed in 2009 by the Indiana Collegiate Action Network (ICAN) and the Indiana Prevention Resource Center (IPRC), with input from Indiana institutions of higher education and the Indiana State Epidemiology and Outcomes Workgroup. The survey was developed to assess substance use and consequences, as well as variables associated with use (availability, peer attitudes and norms, and age of first use). Survey items were adapted from several sources, including the Core Alcohol and Drug Survey (Core Institute, Southern Illinois University - Carbondale), the National College Health Assessment II (American College Health Association), and the Monitoring the Future Survey (Institute for Social Research, University of Michigan). The survey includes three questions on gambling behavior to assess the level and location of gambling among college students and negative consequences experienced by the students. The questions were taken from the Minnesota Student Survey (Stinchfield, Kushner, & Winters, 2005; Winters & Anderson, 2000), and used with the permission of the Minnesota Department of Education.

Several changes were made to the survey instrument in 2012. Two items were added to the survey to assess students' mental health. Additionally, one item was added on sexual orientation, and one item on military service. Response options were changed for the questions on past year and past month substance use. Previously these items provided yes/no and number of days response options. In 2012 the options were changed to number of times, to more closely match the Monitoring the Future Survey. The most extreme response allowed on the two-week binge drinking question was reduced to 14 times, and the most extreme response allowed on the item "How many alcoholic drinks do you think the typical student at your school had the last time he/she "partied/socialized"?" was reduced to 30 drinks in order to eliminate implausibly high responses.

The survey is intended to provide meaningful data at both the campus and state levels, to understand substance use problems and develop effective plans for reducing substance use by college students. Each participating school receives a report of their results. Data from each participating school is the property of that institution, and is not released without their permission. The online survey instrument can be accessed at http://www.drugs.indiana.edu/collegesurvey/?review_year=2012. A copy of the paper version is included at the end of this report.

All two- and four-year colleges in Indiana were invited to participate in the survey. Nine colleges participated in the 2012 survey, including three public and six private schools. Participating schools could choose from either an online or paper format, with all schools but one using the online version. Students were surveyed either before spring break or at least one month after spring break. The period immediately following spring break was avoided because of the historically higher rates of

use during that time. For the online version, students were surveyed during a two-week period. They received an e-mail invitation which included a link to the survey website, and a reminder e-mail midway through the survey period. Incentives were offered by some schools to improve participation rates (e.g., gift cards, athletic tickets, etc.).

Selection of students was determined by the participating school. One school chose to survey a stratified sample of their population, while all others surveyed an entire population of the student body; e.g., all students, all undergraduate students, etc. Participation in the survey was voluntary and anonymous. The survey procedures were reviewed by the Indiana University-Bloomington Institutional Review Board to ensure protection of participants as required by federal Human Subjects regulations.

A total of 47,739 students were invited to participate in the 2012 Indiana College Substance Use Survey, with 7,837 students responding, for a response rate of 16.4%.

Table 2. Response Rate, 2012

Number of students invited to participate	Number of respondents	Response Rate
47,739	7,837	16.4%

Four criteria were used to clean the data. Surveys were discarded if any of the following conditions were met:

- missing age information or inconsistent age responses (indicated 21 years or older but responded to underage questions),
- missing gender information,
- pharmacologically-implausible patterns of responses (i.e., a combination of drugs and frequencies of use whose cumulative effect would be lethal),
- inconsistent responses on substance use items across time measures (monthly vs. annually) for more than one-third of the substances queried.

A total of 869 surveys were rejected due to meeting at least one of the criteria, leaving 6,968 usable surveys (88.9%). Three-fourths of the usable surveys were from students who attended a public institution (5,221), and one-fourth (1,747) from students who attended a private institution.

Table 3. Number of Usable Questionnaires, 2012

	Frequency	Percent
Missing/inconsistent age responses	7	0.1
No gender reported	853	10.9
Implausibly high substance use responses	5	0.1
Inconsistent substance use responses	4	0.1
Total number of usable surveys	6,968	88.9
Total	7,837	100.0

KEY FINDINGS

PREVALENCE AND FREQUENCY OF SUBSTANCE USE

PAST-YEAR PREVALENCE OF USE

More than eighty percent of the students who participated in the 2012 Indiana College Substance Use Survey (81.8%) reported drinking alcohol in the past year, including three-fourths of the students under 21 years of age (74.9%). These rates indicate the pervasiveness of alcohol among the college students participating in the survey. Over one-third of the students (34.0%) reported using marijuana. Almost thirty percent of the students reported smoking tobacco with a hookah (29.4%) or cigarettes (29.2%). Marijuana was the most widely used illicit drug, followed by use of prescription medication not prescribed to the student (12.8%). Annual prevalence rates for all substances investigated are provided in Table 4.

Male students used all of the substances in the past year at significantly higher rates than female students, except for alcohol. Two-fifths of the male students (40.4%) used marijuana in the past year, compared to 30.3% of the female students. Almost half of the male students (43.7%) smoked cigars, and one-third smoked cigarettes (35.7%) or smoked tobacco with a hookah (34.9%). One-fourth of the female students reported smoking cigarettes (25.4%) or tobacco with a hookah (26.2%) in the past year. Approximately one in six male students (15.8%) reported taking prescription medications not prescribed to them, compared to one in nine female students (11.1%). Female students reported a statistically higher rate of alcohol use in the past year (82.5% vs. 80.6% of male students).

Students 21 or older reported higher rates of use in the past year for several substances, including alcohol (89.9%), cigarettes (32.3%), and prescription medications (14.1%), while younger students reported misusing over-the-counter medication at a slightly higher rate (2.6%).

PAST-MONTH PREVALENCE OF USE

Two-thirds of the Indiana students (67.1%) drank alcohol in the past month, including over half of the underage students (58.7%). Almost one in five of the surveyed students (18.7%) used marijuana in the past month, and 15.7% reported cigarette use. Following alcohol, tobacco and marijuana, misuse of prescription drugs was the next most widely used substance by the students in the past month, with 5.3% reporting taking prescription medication not prescribed to them (Table 5).

As with the annual prevalence rates, the male college students reported statistically higher rates of use within the past month for nearly every substance investigated in the survey. One-quarter of the male students (24.3%) reported using marijuana in the past month compared to 15.4% of the female students. Male students used all forms of tobacco at significantly greater rates, with one in five (20.5%) smoking cigarettes, 14.7% smoking cigars, 11.9% using a water pipe, and 9.0% using smokeless tobacco. One in eight of the females (12.9%) smoked cigarettes and 7.3% smoked tobacco with a water pipe. Rates of alcohol use in the past month were not significantly different between the genders.

Students 21 or older reported higher rates of alcohol use (76.9%) and cigarette use (17.1%) in the past month, while younger students reported higher rates of cigar use (8.5%) and smoking tobacco with a hookah (9.9%). Students under 21 also reported a higher rate of over-the-counter drug misuse, with just under one percent (0.9%) reporting the behavior in the past month.

Students attending public institutions were more likely to report use in the past month for almost all of the substances investigated, including alcohol (67.8% vs. 64.9% of students attending private institutions), marijuana (21.8% vs. 9.3%) and cigarettes (17.4% vs. 10.4%). Students attending private institutions reported a higher rate of cigar use in the past month (9.1% vs. 7.1%).

Table 4. Percentage of Students Reporting Use of Select Substances in the Past Year, by Select Demographic Characteristics, 2012

Substance	All Students			Gender		Age		Type of Institution	
	Indiana	U.S. ^a		Males	Females	Under 21	21 or Over	Private	Public
Cigarettes	29.2	25.8		35.7	25.4	26.6	32.3	21.2	31.9
Cigars	25.2			43.7	14.4	25.5	24.8	26.2	24.9
Chewing/smokeless tobacco	7.5			17.1	1.8	7.2	7.9	8.7	7.1
Smoking tobacco with hookah/water pipe	29.4			34.9	26.2	30.2	28.5	24.3	31.1
Alcohol	81.8	77.4		80.6	82.5	74.9	89.9	79.2	82.7
Marijuana	34.0	33.2		40.4	30.3	33.4	34.8	21.3	38.3
Cocaine	3.9	3.3		6.1	2.6	3.0	4.9	1.3	4.7
Hallucinogens	7.0	4.1		10.6	4.9	6.2	8.0	2.3	8.6
Heroin	0.4	0.1		0.9	0.2	0.3	0.5	0.3	0.5
Inhalants	1.1	0.9		1.7	0.7	1.0	1.2	0.7	1.2
Prescription medications not prescribed to student	12.8			15.8	11.1	11.7	14.1	8.1	14.4
Medications prescribed to student, but misused	3.5			4.9	2.7	3.1	4.0	1.9	4.0
Over-the-counter medications misused	2.1			3.1	1.4	2.6	1.4	1.4	2.3
Methamphetamine	0.4	0.2		0.6	0.3	0.3	0.5	0.3	0.4
Other illegal drugs	1.1			1.9	0.7	1.1	1.1	0.8	1.2

^a National data are from the 2011 Monitoring the Future Survey, Johnston et al., 2012.

* $p < .05$, ** $p < .01$, *** $p < .001$

Table 5. Percentage of Students Reporting Use of Select Substances in the Past Month, by Select Demographic Characteristics, 2012

Substance	All Students			Gender		Age			Type of Institution		
	Indiana	U.S. ^a		Males	Females	Under 21	21 or Over		Private	Public	
Cigarettes	15.7	15.2		20.5	12.9	14.5	17.1	**	10.4	17.4	***
Cigars	7.6			14.7	3.5	8.5	6.6	**	9.1	7.1	**
Chewing/smokeless tobacco	3.7			9.0	0.7	3.8	3.7		4.5	3.5	
Smoking tobacco with hookah/water pipe	9.0			11.9	7.3	9.9	8.0	**	7.7	9.4	*
Alcohol	67.1	63.5		66.2	67.6	58.7	76.9	***	64.9	67.8	*
Marijuana	18.7	19.4		24.3	15.4	18.9	18.4		9.3	21.8	***
Cocaine	1.2	1.2		2.1	0.7	1.1	1.3		0.2	1.6	***
Hallucinogens	1.8	1.2		3.0	1.1	2.0	1.6		0.9	2.1	**
Heroin	0.2	+		0.4	0.1	0.1	0.3		0.1	0.2	
Inhalants	0.5	0.3		0.6	0.4	0.5	0.5		0.2	0.6	*
Prescription medications not prescribed to student	5.3			7.2	4.1	4.8	5.8		3.1	6.0	***
Medications prescribed to student, but misused	1.3			1.9	0.9	1.2	1.4		0.8	1.5	*
Over-the-counter medications misused	0.7			1.0	0.5	0.9	0.4	**	0.6	0.7	
Methamphetamine	0.2	+		0.4	0.0	0.2	0.1		0.2	0.2	
Other illegal drugs	0.3			0.7	0.1	0.4	0.2		0.2	0.4	

^a National data are from the 2011 Monitoring the Future Survey, Johnston et al., 2012.. + indicates a prevalence rate of less than 0.05%. **p* < .05, ***p* < .01, ****p* < .001.

PREVALENCE OF BINGE DRINKING

Binge drinking was defined in the Indiana College Substance Use Survey as four or more drinks in one sitting for females and five or more drinks in one sitting for males. This gender-specific standard follows the definition established by the National Institute on Alcohol Abuse and Alcoholism (NIAAA, 2004). As a group, college students are significantly more likely to binge drink than non-college peers (Johnston, O'Malley, Bachman, & Schulenberg, 2009). Students who binge drink experience significantly more adverse consequences than those who do not. One study found that college students who binge drank occasionally were five times more likely to experience five or more alcohol-related problems compared to students who did not binge drink. Students who were frequent binge drinkers (binge drinking three or more times in the past two weeks) were 4 to 15 times more likely to experience a particular problem than non-binge drinkers (Wechsler, Lee, Kuo, & Lee, 2000).

The 2012 Indiana College Substance Use Survey included two questions on binge drinking: in the past month and in the past two weeks. Over half of the students who participated in the 2012 survey (55.1%) reported binge drinking in the past month. Male students reported a higher prevalence rate than female students (59.5% vs. 52.6%), as did students 21 years or older (61.4% vs. 49.7% of underage students), and students at public institutions (56.9%) compared to students attending private institutions (49.7%). See Table 6.

Almost half of the surveyed students (46.0%) reported binge drinking in the past two weeks, with 51.6% of the male students reporting the behavior compared to 42.7% of the female students (Table 7). Over one-fourth of the students (27.8%) reported binge drinking once or twice in the past two weeks, while 18.1% reported doing so three or more times (Figure 1).

Table 6. Percentage of Students Reporting Binge Drinking in the Past Month, by Select Demographic Characteristics, 2012

All Students	Gender			Age		Type of Institution			
	Males	Females		Under 21	21 or Over	Private	Public		
55.1	59.5	52.6	***	49.7	61.4	***	49.7	56.9	***

Note. Binge drinking was defined as consuming 4 or more drinks in one sitting for females and 5 or more drinks in one sitting for males.
 *** $p < .001$

Table 7. Percentage of Students Reporting Binge Drinking in the Past Two Weeks, by Select Demographic Characteristics, 2012


All Students		Gender			Age		Type of Institution			
Indiana	U.S. ^a	Males	Females		Under 21	21 or Over	Private	Public		
46.0	36.1	51.6	42.7	***	41.1	51.6	***	39.9	48.0	***

Note. Binge drinking was defined as consuming 4 or more drinks in one sitting for females and 5 or more drinks in one sitting for males.

^a National data are from the 2011 Monitoring the Future Survey, Johnston et al., 2012.

*** $p < .001$

Figure 1. Frequency of Binge Drinking in the Past Two Weeks, 2012



PRESCRIPTION DRUG MISUSE

According to the Office of National Drug Control Policy, prescription drug abuse is the nation's fastest-growing drug problem (ONDCP, n.d.). College students report higher rates of non-medical use of Adderall and Ritalin than their same-age peers not attending college (Johnston et al., 2012). Among the students participating in the Indiana College Substance Use Survey, Adderall was the most commonly misused prescription medication. Research indicates that the predominant motives reported by college students for misusing prescription stimulants (such as Adderall or Ritalin) are in order to study, improve their concentration, and to stay alert (Teter, McCabe, LaGrange, Cranford, & Boyd, 2006), although nonmedical use of prescription stimulants is associated with lower academic performance (Arria, O'Grady, Caldeira, Vincent, & Wish, 2008; Garnier-Dykstra, Caldeira, Vincent, O'Grady, & Arria, 2012). In a recent study of over 10,000 randomly selected college students from over 100 university, students reporting past year nonmedical prescription stimulant use were ten times more likely to report marijuana use in the past year, almost seven times more likely to report frequent binge drinking, over 20 times more likely to report cocaine use in the past year, and over five times more likely to report driving after binge drinking than college students who had not used prescription stimulants non-medically (McCabe, Knight, Teter, & Wechsler, 2005).

One in ten of the Indiana students (10.5%) reported misusing Adderall in the past six months. Male students and students at public institutions were more likely to misuse Adderall than their counterparts. Approximately one in seven male students (13.8%) reported misusing Adderall in the past six months, and approximately one in nine students attending public institutions (11.9%) did so. Xanax and Vicodin were the next most commonly misused medications, with 3.8% and 3.6% of the students misusing them in the past six months, respectively. Male students misused almost all of the prescription medications at statistically higher rates than female students (Table 8).

Table 8. Percentage of Students Reporting Prescription Drug Misuse in the Past Six Months, by Select Demographic Characteristics, 2012

Substance	All Students	Gender			Age		Type of Institution		
		Males	Females		Under 21	21 or Over	Private	Public	
Adderall	10.5	13.8	8.5	***	10.0	11.0	6.2	11.9	***
Ritalin	1.6	2.7	1.0	***	1.8	1.3	1.4	1.7	
Xanax	3.8	5.2	2.9	***	3.4	4.2	1.7	4.5	***
Codeine	2.8	4.6	1.7	***	3.0	2.6	1.5	3.2	***
Vicodin	3.6	5.4	2.6	***	3.5	3.8	3.3	3.7	
Lortab	1.6	1.7	1.5		0.9	2.4	0.6	1.9	***
Percocet	1.2	1.6	1.0	*	1.2	1.2	1.0	1.2	
Oxycontin	1.0	1.8	0.6	***	0.9	1.2	0.5	1.2	*
Morphine	0.6	1.0	0.4	***	0.6	0.7	0.5	0.7	
Methadone	0.4	0.5	0.3		0.3	0.4	0.2	0.4	
Steroids	0.2	0.4	0.1	*	0.2	0.2	0.1	0.3	


* $p < .05$, *** $p < .001$

COMBINING ALCOHOL WITH ENERGY DRINKS OR OTHER STIMULANTS

College students often cite the desire to counteract the sedative effects of alcohol as a reason for combining it with energy drinks. These students intentionally choose to combine the two in order to be able to drink more alcohol (Marczinski, 2011). Students who mix alcohol and energy drinks consume more alcohol, and experience higher rates of alcohol-related consequences. These include higher prevalence of being taken advantage of sexually, taking advantage of another student sexually, riding in an automobile with a driver under the influence of alcohol, being hurt or injured, and requiring medical treatment (O'Brien, McCoy, Rhodes, Wagoner, & Wolfson, 2008).

Over one-third of the students participating in the Indiana survey who drank alcohol in the past six months (34.9%) reported combining it with energy drinks, and 6.8% combined it with other stimulants, such as Adderall (Figure 2).


Figure 2. Percentage of Students Who Drank Alcohol in Past Six Months, Who Combined Alcohol with Energy Drinks or Other Stimulants, 2012 (n = 5,651)



USUAL TYPE OF ALCOHOL

Over half of the students who drank alcohol in the past six months (53.3%) indicated that liquor was the most common type of alcohol they consumed. One-quarter of the students (25.7%) indicated that beer was the usual type of alcohol they consumed (Figure 3).

Figure 3. Percentage of Students Who Drank Alcohol in Past Six Months, Who Reported Select Categories as the Usual Type of Alcohol They Drank in the Past Month, 2012 (n =5,651)



REASONS FOR ALCOHOL USE

Almost nine in ten of the students who drank alcohol in the past six months (88.8%) reported doing so in order “to have a good time with my friends”. Nearly two-thirds of the students (61.5%) reported drinking alcohol “to relax or relieve tension”, and half (50.1%) did so “to feel good or get high”. Over half of the students (55.7%) said they drink alcohol “because it tastes good”, while one in five students (20.8%) indicated that they used alcohol because of “boredom/nothing else to do” (Table 9).

Male students were more likely to indicate drinking alcohol for most of the reasons investigated, except “to experiment”, which more female than male students indicated (23.3% vs. 19.9%, respectively), and “to get away from my problems” or “because of anger or frustration” where there were no statistical differences between genders.

Students under 21 years old were more likely to report drinking alcohol “to feel good or get high” (52.5%) and “to experiment” (27.4%), while students 21 or older reported higher rates of drinking alcohol “to relax or relieve tension” (67.4%), “because it tastes good” (64.6%), “because of anger or frustration” (16.1%) and “to get to sleep” (12.0%).

Students at public institutions reported higher rates of drinking alcohol for many of the reasons investigated. Almost nine in ten of the students at public institutions who drank alcohol in the past six months (89.4%) indicated that “to have a good time with my friends” was a reason they drink. Over half of the students at public institutions who drink alcohol (52.9%) reported drinking “to feel good or get high”, compared to two-fifths of the students at private institutions who drink (41.5%). Students attending private institutions reported drinking “to experiment” more frequently than those attending public institutions.

Table 9. Percentage of Students Who Drank Alcohol in Past Six Months, Who Indicated Select Reasons for Drinking Alcohol, 2012 (n = 5,651)

Substance	All Students		Gender		Age		Type of Institution				
			Males	Females	Under 21	21 or Over	Private	Public			
To experiment	22.0	19.9	23.3	**	27.4	16.8	***	25.8	20.8	***	
Because I am "hooked"	2.3	3.2	1.8	**	2.1	2.6		1.7		2.6	
To relax or relieve tension	61.5	64.7	59.7	***	55.4	67.4	***	59.7		62.1	
To feel good or get high	50.1	56.6	46.4	***	52.5	47.8	***	41.5		52.9	***
To seek deeper insights and understanding	6.2	9.8	4.2	***	6.2	6.3		5.7		6.4	
To have a good time with my friends	88.8	90.7	87.7	**	88.1	89.4		86.6		89.4	**
To fit in with a group I like	14.5	18.7	12.1	***	14.9	14.0		10.7		15.7	***
Boredom, nothing else to do	20.8	26.7	17.5	***	21.8	19.9		19.8		21.2	
Because of anger or frustration	15.0	15.8	14.6		13.9	16.1	*	14.7		15.1	
To get through the day	3.6	4.9	2.9	***	3.5	3.8		4.0		3.5	
To increase the effects of other drugs	3.9	6.5	2.5	***	4.3	3.6		2.8		4.3	**
To decrease the effects of other drugs	1.6	2.6	1.0	***	1.3	1.9		0.8		1.9	**
To get to sleep	9.7	10.9	9.1	*	7.4	12.0	***	8.1		10.3	*
Because it tastes good	55.7	60.0	53.1	***	46.4	64.6	***	57.1		55.2	
To get away from my problems	13.7	14.0	13.4		13.9	13.4		13.7		13.7	

*p < .05, **p < .01, ***p < .001

CONSEQUENCES OF SUBSTANCE USE

Consequences resulting from alcohol use include violence, unintentional injuries, risky sexual behavior, diminished academic performance, and use of other drugs. Many times the negative consequences from college students' alcohol use affect not only the individuals consuming the alcohol but those around them as well (NIAAA, 2004/2005; Weschler & Nelson, 2008). Researchers estimated that alcohol-related problems experienced by 18-24 year old U.S. college students in 2005 included 1,825 student deaths from alcohol-related injuries and approximately 599,000 unintentional injuries while under the influence of alcohol. The researchers estimated the level of violence associated with alcohol use by college students to be approximately 696,000 students assaulted by other students who had been drinking, including 97,000 sexual assaults (Hingson, Zha, & Weitzman, 2009).

The Indiana College Substance Use Survey asked students to indicate the number of times they experienced select consequences due to their drinking or drug use. Students reported alarming levels of adverse consequences as a result of their drinking. Two-thirds of the students who drank alcohol within the past six months (65.8%) reported having a hangover, and one-third (33.5%) reported blacking out (forgetting where they were or what they did). One in five of the students (19.9%) reported missing class or an assignment as a result of their drinking, and approximately one in seven had driven a car while under the influence (13.5%) and engaged in risky sexual behavior (13.7%) (Table 10).

Approximately one-third of the students who used drugs in the past six months (36.8%) reported driving a car while under the influence, and a similar number (34.3%) reported abusing more than one drug at a time. Sixteen percent of the students who used drugs in the past six months missed class or an assignment due to their drug use, and 13.1% reported having blackouts or flashbacks (Table 11).

Table 10. Percentage of Students Who Drank Alcohol in the Past Six Months, Who Reported Select Consequences as a Result of Their Drinking, 2012 (n = 5,651)

Consequence	%
Friends or family members worried or complained about your drinking	13.8
Had a hangover	65.8
Felt bad or guilty about your drinking	31.3
Been unable to stop drinking when you wanted to	7.8
Created problems between you and your friends or family members	14.0
Gotten into trouble at work or school because of drinking	4.4
Gone to anyone for help to reduce your drinking	2.0
Been hurt or injured because of your drinking	13.1
Been ticketed or arrested because of your drinking	3.3
Been arrested for drunk driving	0.5
Driven a car while under the influence	13.5
Gotten into physical fights when drinking	4.6
Experienced unwanted sexual attention such as being touched when you don't want to be, forced kissing, etc.	11.1
Engaged in risky sexual behavior such as having unprotected sex or sex with someone you just met because you were drinking	13.7
Missed class or an assignment	19.9
Forgot where you were or what you did (blacked out)	33.5

Table 11. Percentage of Students Who Used Drugs in the Past Six Months, Who Reported Select Consequences as a Result of Their Drug Use, 2012 (n = 1,523)

Consequence	%
Abused more than one drug at a time	34.3
Been unable to stop using drugs when you wanted to	8.7
Had blackouts or flashbacks as a result of drug use	13.1
Felt bad or guilty about your drug use	29.2
Had friends or family members worry or complain about your drug use	12.4
Created problems between you and your friends or family members	9.0
Engaged in illegal activities in order to obtain drugs	12.0
Got into physical fights because you were using drugs	1.7
Experienced withdrawal symptoms (felt sick) when you stopped taking drugs	7.9
Had medical problems as a result of your drug use (e.g., memory loss, infections, convulsions, bleeding)	5.4
Driven a car while under the influence of drugs	36.8
Got into trouble at work or school because of your drug use	3.1
Been in vulnerable situations (e.g., could have been hurt and/or date raped when using drugs)	6.3
Engaged in risky sexual behavior such as having unprotected sex or sex with someone you just met because you were using drugs	8.6
Failed a class because of drug use	1.6
Missed class or an assignment due to drug use	16.5

AGE OF FIRST USE

Early onset of substance use is correlated with a range of substance-related problems, including involvement in motor vehicle crashes, unintentional injuries, and alcohol and/or drug dependence (Behrendt, Wittchen, Hofer, Lieb, & Beesdo, 2009; Hingson & Wenxing, 2009). Individuals who begin drinking in their mid-teens are four times more likely to become alcohol dependent during their lifetime compared to those who do not drink before the age of 21 (Grant & Dawson, 1998). The Indiana College Substance Use Survey asked students how old they were when they first used select substances. Responses consisted of two-year intervals for ages 10 through 17 (i.e., 10-11 years, 12-13 years, etc.), a three-year interval for students aged 18 through 20, a four-year interval for ages 21 through 25, and open-ended ranges for extreme responses (“under 10” or “26+”). The mean age of first use was calculated using the midpoints of the intervals and the upper and lower endpoints for the extreme responses.

The mean age of first time use of inhalants was the lowest of all substances investigated, at 16.3 years, followed by over-the-counter medication misuse (16.9 years) and alcohol (17.0 years). Male students initiated use of cigars and alcohol at an earlier average age than female students. Students at public institutions first used all forms of tobacco, and alcohol and marijuana at younger average ages than students at private institutions (Table 12).

Table 12. Average Age of First Time Use of Select Substances, 2012

Substance	All Students	Gender		Type of Institution		
		Males	Females	Private	Public	
Cigarettes	17.1	17.1	17.1	17.5	17.0	***
Cigars	17.7	17.6	17.9	18.1	17.6	***
Chewing/smokeless tobacco	17.3	17.3	17.4	17.7	17.2	**
Smoking tobacco with hookah/water pipe	18.1	18.0	18.1	18.5	18.0	***
Alcohol	17.0	16.9	17.1	17.5	16.9	***
Marijuana	17.4	17.3	17.4	17.8	17.3	***
Cocaine	18.9	19.0	18.7	18.9	18.8	
Hallucinogens	18.6	18.6	18.8	18.9	18.6	
Heroin	18.0	18.0	17.9	18.7	17.9	
Inhalants	16.3	16.4	16.0	16.9	16.2	
Prescription medications not prescribed to student	17.8	17.8	17.9	17.9	17.8	
Medications prescribed to student, but misused	17.5	17.3	17.7	17.6	17.5	
Over-the-counter medications misused	16.9	16.9	16.9	17.3	16.9	
Methamphetamine	17.6	17.9	17.0	18.5	17.4	
Other illegal drugs	17.8	17.9	17.6	18.9	17.6	*

Note. Average age was calculated using the mid-points of the age ranges provided, and the upper and lower endpoints for the extreme responses.

* $p < .05$, ** $p < .01$, *** $p < .001$

ALCOHOL AVAILABILITY

College students' access to alcohol is correlated with increased levels of drinking (Wechsler, Kuo, Lee, & Dowdall, 2000). Students under 21 routinely report that alcohol is easy to obtain from a variety of sources (Brown, Matousek, & Radue, 2009; Fabian, Toomey, Lenk, & Erickson, 2008; Wechsler, Lee, Nelson, & Kuo, 2002). To help understand how underage students obtain alcohol, the Indiana survey asked underage students who drink alcohol to identify the three easiest ways they obtain it. Over half of the students (52.2%) reported that older friends were the easiest way to get alcohol, and almost three-fourths of the students (70.1%) listed older friends as one of the top two ways. "Off-campus private parties" was the most common response for both the second and third easiest method for obtaining alcohol. Over half of the survey respondents (55.6%) selected off-campus private parties as one of the top three easiest ways to access alcohol (Table 13). These data highlight the substantial role that older friends and acquaintances play in how underage students obtain alcohol, and underscore the importance of targeting this group when trying to reduce underage students' access to alcohol.

Students under 21 who drank alcohol in the past year were asked if they had purchased alcohol from a retailer in the past six months, with 10.2% reporting that they had (Figure 4). Of those students, 41.8% reported that purchasing it without being asked for ID was the most common method they used for buying it. Slightly less than one-third of the students who had purchased alcohol from a retailer (31.4%) said that using a fake ID was the most common method they used, while one-quarter of the students (24.7%) indicated that using someone else's ID was the most common method (Figure 5).

Figure 6 shows the locations underage students listed as the easiest place to purchase alcohol, with bars being selected most often (28.2%), followed by restaurants (26.1%).

Table 13. Three Easiest Ways of Getting Alcohol as Reported by Students Who Are Under 21 and Who Drink Alcohol, 2012 (n = 2,827)

	Most common response	2nd most common response	3rd most common response
First Easiest	Older friends (52.2%)	Fraternity or sorority houses (15.5%)	Off-campus private parties (10.6%)
Second Easiest	Off-campus private parties (27.0%)	Older friends (17.9%)	Fraternity or sorority houses (11.7%) On-campus parties (11.7%)
Third Easiest	Off-campus private parties (18.0%)	Someone else on campus (16.0%)	On-campus parties (12.8%)

Note. Underage students who reported alcohol use in the past month or in the past year are included. Students were asked to identify the first, second, and third easiest ways that they obtain alcohol. The top three responses in each category are reported in the table.

Figure 4. Percentage of Students Under 21 Who Drink Alcohol, Who Reported Purchasing Alcohol from a Retailer in the Past Six Months, 2012 (n = 2,827)




Figure 5. Percentage of Students Under 21 Who Have Purchased Alcohol from a Retailer, Who Reported Select Methods as the Most Common for Purchasing Alcohol, 2012 (n = 287)





Figure 6. Percentage of Students Under 21 Who Reported Purchasing Alcohol from Retailers, Easiest Place to Purchase Alcohol, 2012 (n = 287)



PERCEPTIONS OF PEER BEHAVIORS AND BELIEFS

Many college students overestimate both the amount of drinking and drug use, and the approval of use by their peers (Borsari & Carey, 2003; Martens et al., 2006). These misperceptions have been shown to be positively correlated with the student's own drinking and drug use levels. Students who believe their peers drink or use drugs are more likely to drink or use drugs themselves (Kilmer et al., 2006; Neighbors, Lee, Lewis, Fossos, & Larimer, 2007; Perkins, Haines, & Rice, 2005). The Indiana College Substance Use Survey asked students how many alcoholic drinks they thought the typical student at their school had the last time he or she partied/socialized. Students could provide any figure between 0 – 30. The average of the responses was 5.9 drinks (Table 14). The average of the responses is above the threshold for binge drinking, indicating that on average survey respondents thought the typical student binge drank the last time he or she partied. However, the students' responses on their own binge drinking indicate that 46.0% binge drank in the past two weeks. This discrepancy between the percentage of students reporting binge drinking and the average perception of the "typical" student as having binge drank suggests that the students surveyed do indeed overestimate the amount of drinking by their peers. Male students and students under 21 estimated significantly higher numbers of drinks than female and older students.

The survey also asked participants how they thought other students on their campus felt about using certain substances. An astonishingly small 13.7% thought their peers would disapprove (somewhat or strongly) of trying marijuana once or twice, and only one in five (20.9%) thought their peers would disapprove of smoking it occasionally. Less than half (45.4%) thought their peers would disapprove of smoking marijuana regularly. Slightly more than a third of the students (38.6%) thought their peers would disapprove of having five or more drinks in one sitting, and a similar number (35.9%) thought their peers would disapprove of taking one or two drinks nearly every day (Table 15).

Table 14. Mean Number of Drinks Respondents Thought the Typical Student Consumed the Last Time He/She “Partied/Socialized”, 2012 (range = 0 – 30)

	All Students	Gender		Age			Type of Institution	
		Males	Females	Under 21	21 or Over	Private	Public	
Mean	5.9	6.1	5.8 ***	5.8	3.2 **	5.9	5.9	
Standard Deviation	3.2	3.4	3.0	6.0	3.2	2.9	3.3	

** $p < .01$, *** $p < .001$

Table 15. Percentage of Students Who Thought Other Students on Campus Would Approve or Disapprove of Select Behaviors, 2012

	Strongly approve	Somewhat approve	Neither approve or disapprove	Somewhat disapprove	Strongly disapprove	No answer
Trying marijuana once or twice	22.6	38.0	25.3	6.6	7.1	0.4
Smoking marijuana occasionally	13.7	35.2	29.6	11.9	9.0	0.6
Smoking marijuana regularly	5.9	19.1	29.0	28.3	17.1	0.6
Trying cocaine once or twice	1.6	8.1	16.0	35.8	37.9	0.6
Taking cocaine regularly	0.9	1.9	9.2	18.2	68.9	0.8
Trying amphetamines (e.g., Adderall, Ritalin) once or twice	10.3	23.8	27.3	16.8	21.3	0.5
Taking amphetamines regularly	3.9	10.7	21.7	24.1	38.5	1.0
Having one or two alcoholic drinks nearly every day	10.5	22.6	30.5	24.1	11.8	0.6
Having three or four alcoholic drinks nearly every day	4.5	11.6	22.0	30.7	30.5	0.6
Having five or more alcoholic drinks in one sitting	12.2	21.0	27.4	17.9	20.7	0.7
Taking steroids for body building or improved athletic performance	1.5	6.1	23.3	29.4	39.0	0.7

GAMBLING BEHAVIORS

Opportunities for gambling have increased rapidly since the 1990's, both nationally and in Indiana. In 1989 the State of Indiana began operating the Hoosier Lottery. River boat casinos and horse tracks were approved by the Indiana legislature in the mid-1990s, and in 2007 the State allowed slot machines to be placed in horse parks. The advent of online gambling and the rapid growth in the popularity of poker have also added to the current environment of gambling. A recent national survey of 18-21 year olds (Barnes, Welge, Hoffman & Tidwell, 2010) found that 75% of the college students reported gambling in the past year, which was similar to the rate reported by the non-college young adults surveyed. The rate of frequent gambling (weekly or more) was lower for U.S. college students (18%) than their non-college peers (25%). Male college students reported significantly higher prevalence rates of gambling (83% vs. 67%) and frequent gambling (31% vs. 6%) than the female students.

The most common form of gambling reported by students participating in the 2012 Indiana College Substance Use Survey was playing the lottery (24.6%), followed by playing cards at home (15.2%). Male Indiana students reported higher rates of gambling than female students for all gambling behaviors investigated. Over one-fourth of the male students (26.6%) played cards at home for money in the past year, compared to 8.6% of the female students. Almost one-fourth of the male students (24.0%) gambled in betting pools and nearly one in five (18.0%) gambled at a casino. Students 21 or older reported significantly higher rates than the younger students for most gambling behaviors, with approximately one-fourth of the older students playing the lottery (27.6%) and gambling at a casino (23.6%) (Table 16).

Students were asked how frequently they gambled to assess the level of excessive gambling. Less than one percent of the students reported participating in any of the gambling behaviors at least once a week in the past year (Figure 7). The most frequent location that students reported for gambling (other than online or playing the lottery) was at a casino (9.3%), followed by an apartment or house (6.9%) (Figure 8). The most common negative consequence of gambling reported by Indiana students was feeling bad about their gambling (3.8%), followed by financial issues (1.7%) (Table 17).

Table 16. Percentage of Students Reporting Gambling Behaviors in the Past Year, 2012

Gambling Behavior	All Students		Gender		Age		Type of Institution	
			Males	Females	Under 21	21 or Over	Private	Public
Played cards at home	15.2		26.6	8.6 ***	15.5	14.9	15.4	15.2
Gambled at a casino	13.1		18.0	10.3 ***	4.1	23.6 ***	13.6	12.9
Gambled on-line	2.0		4.5	0.6 ***	1.6	2.5 *	1.5	2.2
Gambled at informal locations (e.g., street corners, gambling houses)	1.3		3.0	0.4 ***	1.2	1.6	1.1	1.4
Bet on fantasy football/basketball	8.4		18.7	2.3 ***	7.4	9.5 **	7.7	8.6
Betting pools (e.g., March Madness)	11.9		24.0	4.8 ***	11.0	13.0 *	13.4	11.4 *
Lottery	24.6		28.1	22.5 ***	22.1	27.6 ***	26.7	23.9 *
Placed bet with bookie	1.5		3.2	0.5 ***	1.4	1.7	1.0	1.7

* $p < .05$, ** $p < .01$, *** $p < .001$

Figure 7. Percentage of Students Who Reported Weekly or More Frequent Gambling in the Past Year, by Gambling Category, 2012




Figure 8. Most Frequent Gambling Location (Other Than Online or Lottery) in Past Year, 2012




Table 17. Percentage of Students Who Gambled in the Past Year, Who Reported Select Consequences as a Result of Their Gambling, 2012 (n = 2,811)

Consequence	%
Academic problems (missed class, did poorly on exam, lower grades)	0.2
Issues with legal system	0.2
Financial issues	1.7
Felt bad about gambling	3.8
Poor physical health (sleep issues, depression, poor nutrition, poor hygiene)	0.4
Issues with friends and family (lied, strained relationships, arguments)	0.9

CHARACTERISTICS OF SURVEY PARTICIPANTS

Almost two-thirds of the survey respondents are female (63.1%) and slightly more than half (53.9%) are less than 21 years of age. Over four-fifths of the students (84.4%) are White and most (95.5%) are non-Hispanic. Respondents were nearly evenly divided among freshmen, sophomores, juniors, and seniors. Over ninety percent (97.0%) were full-time students and slightly more than half (54.8%) lived on-campus, with almost two-fifths (38.8%) living in residence halls. Three-fourths of the students (74.9%) attend public institutions.

Table 18. Demographic Characteristics of Survey Participants, 2012

	Frequency	%
Gender		
Male	2,572	36.9
Female	4,396	63.1
Age		
Under 21	3,755	53.9
21 or older	3,213	46.1
Race		
White	5,879	84.4
Black/African American	239	3.4
Asian	502	7.2
Native American/Alaskan Native	28	0.4
Hawaiian/Pacific Islander	9	0.1
More than one race	163	2.3
Other	103	1.5
No answer	45	0.6
Ethnicity		
Hispanic	259	3.7
Non-Hispanic	6,654	95.5
No answer	55	0.8
Classification		
Freshman	1,794	25.7
Sophomore	1,677	24.1
Junior	1,655	23.8
Senior	1,645	23.6
Grad/professional	108	1.5
Not seeking a degree	27	0.4
Other	41	0.6
No answer	21	0.3

Table 18 (continued). Demographic characteristics of survey participants

	Frequency	%
Student status		
Full-time	6,758	97.0
Part-time	169	2.4
No answer	41	0.6
Location of residence		
On-campus	3,816	54.8
Off-campus	3,106	44.6
No answer	46	0.7
Type of residence		
House/apartment/etc.	3,393	48.7
Residence hall	2,707	38.8
Fraternity/Sorority	530	7.6
Other campus housing	176	2.5
Other	125	1.8
No answer	37	0.5
Type of institution		
Private	1,747	25.1
Public	5,221	74.9

REFERENCES

- Arria, A. M., O'Grady, K. E., Caldeira, K. M., Vincent, K. B., & Wish, E. D. (2008). Nonmedical use of prescription stimulants and analgesics: Associations with social and academic behaviors among college students. *Journal Of Drug Issues, 38*(4), 1045-1060.
- Barnes, G.M., Welge, J.W., Hoffman, J.H., & Tidwell, M.O. (2010) Comparisons of gambling and alcohol use among college students and non-college young people in the United States. *Journal of American College Health, 58*(5), 443-452.
- Behrendt, S., Wittchen, H. U., Höfler, M., Lieb, R., & Beesdo, K. (2009). Transitions from first substance use to substance use disorders in adolescence: Is early onset associated with a rapid escalation? *Drug and Alcohol Dependence, 99*(1-3), 68-78.
- Borsari, B., & Carey, K. B. (2003). Descriptive and injunctive norms in college drinking: A meta-analytic integration. *Journal of Studies on Alcohol, 64*(3), 331.
- Brown, R. L., Matousek, T. A., & Radue, M. B. (2009). Legal-age students' provision of alcohol to underage college students: An exploratory study. *Journal of American College Health, 57*(6), 611-618.
- Fabian, L. E. A., Toomey, T. L., Lenk, K. M., & Erickson, D. J. (2008). How do underage college students get alcohol? *Journal of Drug Education, 38*(1), 15-26.
- Garnier-Dykstra, L. M., Caldeira, K. M., Vincent, K. B., O'Grady, K. E., & Arria, A. M. (2012). Nonmedical use of prescription stimulants during college: Four-year trends in exposure opportunity, use, motives, and sources. *Journal Of American College Health, 60*(3), 226-234.
- Grant, B. F., & Dawson, D. A. (1998). Age of onset of drug use and its association with DSM-IV drug abuse and dependence: results from the National Longitudinal Alcohol Epidemiologic Survey. *Journal Of Substance Abuse, 10*(2), 163-173.
- Hingson, R.W., & Wenxing, Z. (2009). Age of drinking onset, alcohol use disorders, frequent heavy drinking, and unintentionally injuring oneself and others after drinking. *Pediatrics, 123*(6), 1477-1484.
- Hingson, R.W., Zha, W., & Weitzman, E.R. (2009). Magnitude of and trends in alcohol-related mortality and morbidity among U.S. college students age 18-24: 1998 to 2005. *Journal of Studies on Alcohol and Drugs Supplement, 16*, 12-20.
- Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J.E. (2009). *Monitoring the Future, National Survey Results on Drug Use, 1975-2008. Volume II: College Students and Adults Ages 19-50*. NIH Pub. No. 09-7403. Bethesda, MD: National Institute on Drug Abuse.
- Johnston, L.D., O'Malley, P.M., Bachman, J.G., & Schulenberg, J. E. (2012). *Monitoring the Future National Survey Results on Drug Use, 1975-2011. Volume II: College Students and Adults Ages 19-50*. Ann Arbor: Institute for Social Research, The University of Michigan.
- Kilmer, J. R., Walker, D. D., Lee, C. M., Palmer, R. S., Mallett, K. A., Fabiano, P., et al. (2006). Misperceptions of college student marijuana use: Implications for prevention. *Journal of Studies on Alcohol, 67*(2), 277-281.

- Marczinski C.A. (2011). Alcohol mixed with energy drinks: consumption patterns and motivations for use in U.S. college students. *International Journal of Environmental Research And Public Health*, 8(8), 3232-3245.
- Martens, M. P., Page, J. C., Mowry, E. S., Damann, K. M., Taylor, K. K., & Cimini, M. D. (2006). Differences between actual and perceived student norms: An examination of alcohol use, drug use, and sexual behavior. *Journal of American College Health*, 54(5), 295-300.
- McCabe, S., Knight, J. R., Teter, C. J., & Wechsler, H. (2005). Non-medical use of prescription stimulants among US college students: Prevalence and correlates from a national survey. *Addiction*, 100(1), 96-106.
- National Institute on Alcohol Abuse and Alcoholism, Interdisciplinary Team on Underage Drinking Research. (2004/2005). The scope of the problem. *Alcohol Research & Health*, 28(3), 111-120.
- National Institute on Alcohol Abuse and Alcoholism. (2002). *A call to action: Changing the culture of drinking at U.S. colleges* (NIH Pub. No. 02-5010). Bethesda, MD: NIAAA.
- National Institute on Alcohol Abuse and Alcoholism. (2004). NIAAA council approves definition of binge drinking. *NIAAA Newsletter* (3), 3. Retrieved from http://pubs.niaaa.nih.gov/publications/Newsletter/winter2004/Newsletter_Number3.pdf
- Neighbors, C., Lee, C. M., Lewis, M. A., Fossos, N., & Larimer, M. E. (2007). Are social norms the best predictor of outcomes among heavy-drinking college students? *Journal of Studies on Alcohol & Drugs*, 68(4), 556-565.
- O'Brien, M., McCoy, T., Rhodes, S., Wagoner, A., & Wolfson, M. (2008). Caffeinated cocktails: Energy drink consumption, high-risk drinking, and alcohol-related consequences among college students. *Academic Emergency Medicine: Official Journal Of The Society For Academic Emergency Medicine*, 15(5), 453-460.
- Office of National Drug Control Policy (n.d.). *Epidemic: Responding to America's prescription drug abuse crisis*. Retrieved from <http://www.whitehouse.gov/ondcp/prescription-drug-abuse>.
- Perkins, H. W., Haines, M. P., & Rice, R. (2005). Misperceiving the college drinking norm and related problems: A nationwide study of exposure to prevention information, perceived norms and student alcohol misuse. *Journal of Studies on Alcohol*, 470-478.
- Stinchfield, R., Kushner, M. G., & Winters, K. C. (2005). Alcohol use and prior substance abuse treatment in relation to gambling problem severity and gambling treatment outcome. *Journal of Gambling Studies*, 21(3), 273-297.
- Teter, C. J., McCabe, S., LaGrange, K., Cranford, J. A., & Boyd, C. J. (2006). Illicit use of specific prescription stimulants among college students: Prevalence, motives, and routes of administration. *Pharmacotherapy*, 26(10), 1501-1510.
- Wechsler, H., & Nelson, T. F. (2008). What we have learned from the Harvard School of Public Health College Alcohol Study: Focusing attention on college student alcohol consumption and the environmental conditions that promote it. *Journal of Studies on Alcohol & Drugs*, 69(4), 481-490.

- Wechsler, H., Kuo, M., Lee, H., & Dowdall, G. W. (2000). Environmental correlates of underage alcohol use and related problems of college students. *American Journal Of Preventive Medicine, 19*(1), 24-29.
- Wechsler, H., Lee, J. E., Kuo, M., & Lee, H. (2000). College binge drinking in the 1990s: A continuing problem. *Journal of American College Health, 48*(5), 199.
- Wechsler, H., Lee, J. E., Nelson, T. F., & Kuo, M. (2002). Underage college students' drinking behavior, access to alcohol, and the influence of deterrence policies. *Journal of American College Health, 50*(5), 223.
- Winters, K. C., & Anderson, N. (2000). Gambling involvement and drug use among adolescents. *Journal Of Gambling Studies, 16*(2-3), 175-198.

FREQUENCY TABLES

Substance use during the last 12 months: Cigarettes

	Frequency	Percent	Valid Percent	Cumulative Percent
No answer	9	.1	.1	.1
Never	4924	70.7	70.7	70.8
1-5 times	921	13.2	13.2	84.0
Valid 6-19 times	304	4.4	4.4	88.4
20-39 times	178	2.6	2.6	90.9
40+ times	632	9.1	9.1	100.0
Total	6968	100.0	100.0	

Substance use during the last 12 months: Cigars

	Frequency	Percent	Valid Percent	Cumulative Percent
No answer	25	.4	.4	.4
Never	5188	74.5	74.5	74.8
1-5 times	1245	17.9	17.9	92.7
Valid 6-19 times	365	5.2	5.2	97.9
20-39 times	88	1.3	1.3	99.2
40+ times	57	.8	.8	100.0
Total	6968	100.0	100.0	

Substance use during the last 12 months: Chewing or smokeless tobacco

	Frequency	Percent	Valid Percent	Cumulative Percent
No answer	46	.7	.7	.7
Never	6400	91.8	91.8	92.5
1-5 times	250	3.6	3.6	96.1
Valid 6-19 times	82	1.2	1.2	97.3
20-39 times	41	.6	.6	97.9
40+ times	149	2.1	2.1	100.0
Total	6968	100.0	100.0	

Substance use during the last 12 months: Smoking tobacco with hookah or water pipe

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	53	.8	.8	.8
	Never	4865	69.8	69.8	70.6
	1-5 times	1308	18.8	18.8	89.4
	6-19 times	478	6.9	6.9	96.2
	20-39 times	164	2.4	2.4	98.6
	40+ times	100	1.4	1.4	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	32	.5	.5	.5
	Never	1237	17.8	17.8	18.2
	1-5 times	1066	15.3	15.3	33.5
	6-19 times	1239	17.8	17.8	51.3
	20-39 times	1192	17.1	17.1	68.4
	40+ times	2202	31.6	31.6	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Marijuana

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	35	.5	.5	.5
	Never	4559	65.4	65.4	65.9
	1-5 times	1024	14.7	14.7	80.6
	6-19 times	399	5.7	5.7	86.4
	20-39 times	238	3.4	3.4	89.8
	40+ times	713	10.2	10.2	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Cocaine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	25	.4	.4	.4
	Never	6673	95.8	95.8	96.1
	1-5 times	200	2.9	2.9	99.0
	6-19 times	48	.7	.7	99.7
	20-39 times	11	.2	.2	99.8
	40+ times	11	.2	.2	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Hallucinogens

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	23	.3	.3	.3
	Never	6455	92.6	92.6	93.0
	1-5 times	390	5.6	5.6	98.6
	6-19 times	79	1.1	1.1	99.7
	20-39 times	15	.2	.2	99.9
	40+ times	6	.1	.1	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Heroin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	29	.4	.4	.4
	Never	6909	99.2	99.2	99.6
	1-5 times	9	.1	.1	99.7
	6-19 times	11	.2	.2	99.9
	20-39 times	3	.0	.0	99.9
	40+ times	7	.1	.1	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Inhalants

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.4	.4	.4
	Never	6867	98.6	98.6	98.9
	1-5 times	51	.7	.7	99.7
	6-19 times	14	.2	.2	99.9
	20-39 times	4	.1	.1	99.9
	40+ times	5	.1	.1	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Prescription pills that are not prescribed to you

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	31	.4	.4	.4
	Never	6043	86.7	86.7	87.2
	1-5 times	523	7.5	7.5	94.7
	6-19 times	235	3.4	3.4	98.0
	20-39 times	76	1.1	1.1	99.1
	40+ times	60	.9	.9	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Medications prescribed to you but used to get high

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.4	.4	.4
	Never	6697	96.1	96.1	96.5
	1-5 times	142	2.0	2.0	98.5
	6-19 times	57	.8	.8	99.4
	20-39 times	26	.4	.4	99.7
	40+ times	19	.3	.3	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Over-the-counter medications to get high

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	58	.8	.8	.8
	Never	6767	97.1	97.1	97.9
	1-5 times	106	1.5	1.5	99.5
	6-19 times	29	.4	.4	99.9
	20-39 times	6	.1	.1	100.0
	40+ times	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Methamphetamine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	30	.4	.4	.4
	Never	6911	99.2	99.2	99.6
	1-5 times	14	.2	.2	99.8
	6-19 times	3	.0	.0	99.9
	20-39 times	6	.1	.1	99.9
	40+ times	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Substance use during the last 12 months: Other illegal drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	759	10.9	10.9	10.9
	Never	6131	88.0	88.0	98.9
	1-5 times	61	.9	.9	99.8
	6-19 times	7	.1	.1	99.9
	20-39 times	4	.1	.1	99.9
	40+ times	6	.1	.1	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Cigarettes

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	370	5.3	5.3	5.3
	Never	5506	79.0	79.0	84.3
	1-5 times	520	7.5	7.5	91.8
	6-19 times	165	2.4	2.4	94.2
	20-39 times	119	1.7	1.7	95.9
	40+ times	288	4.1	4.1	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Cigars

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	389	5.6	5.6	5.6
	Never	6047	86.8	86.8	92.4
	1-5 times	486	7.0	7.0	99.3
	6-19 times	29	.4	.4	99.8
	20-39 times	8	.1	.1	99.9
	40+ times	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Chewing or smokeless tobacco

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	409	5.9	5.9	5.9
	Never	6299	90.4	90.4	96.3
	1-5 times	113	1.6	1.6	97.9
	6-19 times	53	.8	.8	98.7
	20-39 times	40	.6	.6	99.2
	40+ times	54	.8	.8	100.0
	Total	6968	100.0	100.0	

**Number of times in the past month substance was used: Smoking tobacco
with hookah or water pipe**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	410	5.9	5.9	5.9
	Never	5932	85.1	85.1	91.0
	1-5 times	547	7.9	7.9	98.9
	6-19 times	59	.8	.8	99.7
	20-39 times	15	.2	.2	99.9
	40+ times	5	.1	.1	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	386	5.5	5.5	5.5
	Never	1907	27.4	27.4	32.9
	1-5 times	2522	36.2	36.2	69.1
	6-19 times	1761	25.3	25.3	94.4
	20-39 times	267	3.8	3.8	98.2
	40+ times	125	1.8	1.8	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Marijuana

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	392	5.6	5.6	5.6
	Never	5274	75.7	75.7	81.3
	1-5 times	666	9.6	9.6	90.9
	6-19 times	297	4.3	4.3	95.1
	20-39 times	166	2.4	2.4	97.5
	40+ times	173	2.5	2.5	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Cocaine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	384	5.5	5.5	5.5
	Never	6499	93.3	93.3	98.8
	1-5 times	78	1.1	1.1	99.9
	6-19 times	3	.0	.0	99.9
	20-39 times	2	.0	.0	100.0
	40+ times	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Hallucinogens

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	382	5.5	5.5	5.5
	Never	6460	92.7	92.7	98.2
	1-5 times	119	1.7	1.7	99.9
	6-19 times	4	.1	.1	100.0
	40+ times	3	.0	.0	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Heroin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	392	5.6	5.6	5.6
	Never	6563	94.2	94.2	99.8
	1-5 times	9	.1	.1	99.9
	6-19 times	3	.0	.0	100.0
	40+ times	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Inhalants

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	388	5.6	5.6	5.6
	Never	6548	94.0	94.0	99.5
	1-5 times	26	.4	.4	99.9
	6-19 times	5	.1	.1	100.0
	20-39 times	1	.0	.0	100.0
	Total	6968	100.0	100.0	

**Number of times in the past month substance was used: Prescription pills
that are not prescribed to you**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	387	5.6	5.6	5.6
	Never	6215	89.2	89.2	94.7
	1-5 times	286	4.1	4.1	98.9
	6-19 times	66	.9	.9	99.8
	20-39 times	9	.1	.1	99.9
	40+ times	5	.1	.1	100.0
	Total	6968	100.0	100.0	

**Number of times in the past month substance was used: Medications
prescribed to you but used to get high**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	384	5.5	5.5	5.5
	Never	6494	93.2	93.2	98.7
	1-5 times	68	1.0	1.0	99.7
	6-19 times	16	.2	.2	99.9
	20-39 times	5	.1	.1	100.0
	40+ times	1	.0	.0	100.0
	Total	6968	100.0	100.0	

**Number of times in the past month substance was used: Over-the-counter
medications to get high**

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	404	5.8	5.8	5.8
	Never	6516	93.5	93.5	99.3
	1-5 times	38	.5	.5	99.9
	6-19 times	10	.1	.1	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Methamphetamine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	417	6.0	6.0	6.0
	Never	6540	93.9	93.9	99.8
	1-5 times	5	.1	.1	99.9
	6-19 times	3	.0	.0	100.0
	20-39 times	1	.0	.0	100.0
	40+ times	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Number of times in the past month substance was used: Other illegal drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	457	6.6	6.6	6.6
	Never	6488	93.1	93.1	99.7
	1-5 times	18	.3	.3	99.9
	6-19 times	4	.1	.1	100.0
	40+ times	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Number of days student binge drank during the past month

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	32	.5	.5	.5
	0	3095	44.4	44.4	44.9
	1	882	12.7	12.7	57.5
	2	655	9.4	9.4	66.9
	3	432	6.2	6.2	73.1
	4	375	5.4	5.4	78.5
	5	334	4.8	4.8	83.3
	6	211	3.0	3.0	86.3
	7	143	2.1	2.1	88.4
	8	213	3.1	3.1	91.4
	9	64	.9	.9	92.4
	10	220	3.2	3.2	95.5
	11	29	.4	.4	95.9
	12	107	1.5	1.5	97.5
	13	21	.3	.3	97.8
	14	22	.3	.3	98.1
	15	53	.8	.8	98.9
	16	15	.2	.2	99.1
	17	7	.1	.1	99.2
	18	9	.1	.1	99.3
	20	23	.3	.3	99.6
	21	4	.1	.1	99.7
	22	3	.0	.0	99.7
	23	1	.0	.0	99.7
	24	3	.0	.0	99.8
	25	4	.1	.1	99.8
	26	2	.0	.0	99.9
	29	1	.0	.0	99.9
	30	8	.1	.1	100.0
	Total	6968	100.0	100.0	

Number of times student binge drank during the last two weeks

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	37	.5	.5	.5
	0	3727	53.5	53.5	54.0
	1	1162	16.7	16.7	70.7
	2	772	11.1	11.1	81.8
	3	407	5.8	5.8	87.6
	4	345	5.0	5.0	92.6
	5	212	3.0	3.0	95.6
	6	145	2.1	2.1	97.7
	7	44	.6	.6	98.3
	8	40	.6	.6	98.9
	9	13	.2	.2	99.1
	10	26	.4	.4	99.5
	11	4	.1	.1	99.5
	12	14	.2	.2	99.7
	13	2	.0	.0	99.7
	14	18	.3	.3	100.0
	Total	6968	100.0	100.0	

Age of first time use: Cigarettes

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	9	.1	.1	.1
	Did not use	4264	61.2	61.2	61.3
	Under 10	56	.8	.8	62.1
	10-11	56	.8	.8	62.9
	12-13	166	2.4	2.4	65.3
	14-15	344	4.9	4.9	70.2
	16-17	785	11.3	11.3	81.5
	18-20	1167	16.7	16.7	98.3
	21-25	121	1.7	1.7	100.0
	Total	6968	100.0	100.0	

Age of first time use: Cigars

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	32	.5	.5	.5
	Did not use	4516	64.8	64.8	65.3
	Under 10	13	.2	.2	65.5
	10-11	13	.2	.2	65.6
	12-13	51	.7	.7	66.4
	14-15	241	3.5	3.5	69.8
	16-17	788	11.3	11.3	81.1
	18-20	1214	17.4	17.4	98.6
	21-25	91	1.3	1.3	99.9
	26+	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Chewing or smokeless tobacco

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	49	.7	.7	.7
	Did not use	6049	86.8	86.8	87.5
	Under 10	13	.2	.2	87.7
	10-11	9	.1	.1	87.8
	12-13	31	.4	.4	88.3
	14-15	109	1.6	1.6	89.8
	16-17	286	4.1	4.1	93.9
	18-20	387	5.6	5.6	99.5
	21-25	34	.5	.5	100.0
	26+	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Smoking tobacco with hookah or water pipe

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	41	.6	.6	.6
	Did not use	4376	62.8	62.8	63.4
	Under 10	5	.1	.1	63.5
	10-11	5	.1	.1	63.5
	12-13	10	.1	.1	63.7
	14-15	157	2.3	2.3	65.9
	16-17	770	11.1	11.1	77.0
	18-20	1506	21.6	21.6	98.6
	21-25	93	1.3	1.3	99.9
	26+	5	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	20	.3	.3	.3
	Did not use	1026	14.7	14.7	15.0
	Under 10	106	1.5	1.5	16.5
	10-11	60	.9	.9	17.4
	12-13	300	4.3	4.3	21.7
	14-15	1095	15.7	15.7	37.4
	16-17	1826	26.2	26.2	63.6
	18-20	2204	31.6	31.6	95.2
	21-25	327	4.7	4.7	99.9
	26+	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Marijuana

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	30	.4	.4	.4
	Did not use	3979	57.1	57.1	57.5
	Under 10	8	.1	.1	57.6
	10-11	16	.2	.2	57.9
	12-13	91	1.3	1.3	59.2
	14-15	453	6.5	6.5	65.7
	16-17	984	14.1	14.1	79.8
	18-20	1283	18.4	18.4	98.2
	21-25	117	1.7	1.7	99.9
	26+	7	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Cocaine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	23	.3	.3	.3
	Did not use	6550	94.0	94.0	94.3
	Under 10	6	.1	.1	94.4
	12-13	2	.0	.0	94.4
	14-15	18	.3	.3	94.7
	16-17	76	1.1	1.1	95.8
	18-20	226	3.2	3.2	99.0
	21-25	63	.9	.9	99.9
	26+	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Hallucinogens

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Did not use	6270	90.0	90.0	90.3
	Under 10	7	.1	.1	90.4
	12-13	3	.0	.0	90.4
	14-15	22	.3	.3	90.7
	16-17	154	2.2	2.2	92.9
	18-20	415	6.0	6.0	98.9
	21-25	75	1.1	1.1	100.0
	26+	3	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Heroin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	30	.4	.4	.4
	Did not use	6877	98.7	98.7	99.1
	Under 10	7	.1	.1	99.2
	14-15	4	.1	.1	99.3
	16-17	9	.1	.1	99.4
	18-20	32	.5	.5	99.9
	21-25	7	.1	.1	100.0
	26+	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Inhalants

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	25	.4	.4	.4
	Did not use	6787	97.4	97.4	97.8
	Under 10	10	.1	.1	97.9
	10-11	6	.1	.1	98.0
	12-13	13	.2	.2	98.2
	14-15	32	.5	.5	98.6
	16-17	43	.6	.6	99.3
	18-20	40	.6	.6	99.8
	21-25	12	.2	.2	100.0
	Total	6968	100.0	100.0	

Age of first time use: Prescription pills that are not prescribed to you

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	17	.2	.2	.2
	Did not use	5904	84.7	84.7	85.0
	Under 10	11	.2	.2	85.1
	10-11	6	.1	.1	85.2
	12-13	21	.3	.3	85.5
	14-15	121	1.7	1.7	87.3
	16-17	302	4.3	4.3	91.6
	18-20	501	7.2	7.2	98.8
	21-25	76	1.1	1.1	99.9
	26+	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Age of first time use: Medications prescribed to you but used to get high

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	29	.4	.4	.4
	Did not use	6558	94.1	94.1	94.5
	Under 10	6	.1	.1	94.6
	10-11	3	.0	.0	94.7
	12-13	8	.1	.1	94.8
	14-15	46	.7	.7	95.4
	16-17	142	2.0	2.0	97.5
	18-20	148	2.1	2.1	99.6
	21-25	26	.4	.4	100.0
	26+	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Over-the-counter medications to get high

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	34	.5	.5	.5
	Did not use	6686	96.0	96.0	96.4
	Under 10	6	.1	.1	96.5
	10-11	2	.0	.0	96.6
	12-13	9	.1	.1	96.7
	14-15	53	.8	.8	97.4
	16-17	78	1.1	1.1	98.6
	18-20	89	1.3	1.3	99.8
	21-25	10	.1	.1	100.0
	26+	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Methamphetamine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	26	.4	.4	.4
	Did not use	6879	98.7	98.7	99.1
	Under 10	7	.1	.1	99.2
	14-15	5	.1	.1	99.3
	16-17	19	.3	.3	99.5
	18-20	22	.3	.3	99.9
	21-25	9	.1	.1	100.0
	26+	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Age of first time use: Other illegal drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	58	.8	.8	.8
	Did not use	6795	97.5	97.5	98.3
	Under 10	7	.1	.1	98.5
	12-13	1	.0	.0	98.5
	14-15	12	.2	.2	98.6
	16-17	26	.4	.4	99.0
	18-20	57	.8	.8	99.8
	21-25	12	.2	.2	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Adderall

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	29	.4	.4	.4
	Did not use	6209	89.1	89.1	89.5
	Once	301	4.3	4.3	93.8
	Every other month	146	2.1	2.1	95.9
	Once a month	153	2.2	2.2	98.1
	Once a week	83	1.2	1.2	99.3
	3 times a week	32	.5	.5	99.8
	5 times a week	7	.1	.1	99.9
	Every day	8	.1	.1	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Ritalin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	41	.6	.6	.6
	Did not use	6815	97.8	97.8	98.4
	Once	58	.8	.8	99.2
	Every other month	10	.1	.1	99.4
	Once a month	23	.3	.3	99.7
	Once a week	12	.2	.2	99.9
	3 times a week	3	.0	.0	99.9
	5 times a week	2	.0	.0	99.9
	Every day	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Xanax

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	40	.6	.6	.6
	Did not use	6665	95.7	95.7	96.2
	Once	130	1.9	1.9	98.1
	Every other month	45	.6	.6	98.7
	Once a month	53	.8	.8	99.5
	Once a week	18	.3	.3	99.8
	3 times a week	11	.2	.2	99.9
	5 times a week	2	.0	.0	99.9
	Every day	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Codeine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	39	.6	.6	.6
	Did not use	6735	96.7	96.7	97.2
	Once	100	1.4	1.4	98.7
	Every other month	43	.6	.6	99.3
	Once a month	41	.6	.6	99.9
	Once a week	6	.1	.1	99.9
	3 times a week	2	.0	.0	100.0
	5 times a week	1	.0	.0	100.0
	Every day	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Vicodin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	45	.6	.6	.6
	Did not use	6670	95.7	95.7	96.4
	Once	136	2.0	2.0	98.3
	Every other month	50	.7	.7	99.0
	Once a month	35	.5	.5	99.5
	Once a week	16	.2	.2	99.8
	3 times a week	9	.1	.1	99.9
	5 times a week	4	.1	.1	100.0
	Every day	3	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Lortab

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	44	.6	.6	.6
	Did not use	6813	97.8	97.8	98.4
	Once	61	.9	.9	99.3
	Every other month	15	.2	.2	99.5
	Once a month	17	.2	.2	99.7
	Once a week	10	.1	.1	99.9
	3 times a week	2	.0	.0	99.9
	5 times a week	2	.0	.0	99.9
	Every day	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Percocet

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	47	.7	.7	.7
	Did not use	6838	98.1	98.1	98.8
	Once	36	.5	.5	99.3
	Every other month	21	.3	.3	99.6
	Once a month	15	.2	.2	99.8
	Once a week	6	.1	.1	99.9
	3 times a week	2	.0	.0	100.0
	5 times a week	1	.0	.0	100.0
	Every day	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Oxycontin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	47	.7	.7	.7
	Did not use	6850	98.3	98.3	99.0
	Once	42	.6	.6	99.6
	Every other month	8	.1	.1	99.7
	Once a month	7	.1	.1	99.8
	Once a week	9	.1	.1	99.9
	3 times a week	2	.0	.0	100.0
	5 times a week	1	.0	.0	100.0
	Every day	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Morphine

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	43	.6	.6	.6
	Did not use	6882	98.8	98.8	99.4
	Once	23	.3	.3	99.7
	Every other month	5	.1	.1	99.8
	Once a month	5	.1	.1	99.9
	Once a week	8	.1	.1	100.0
	3 times a week	1	.0	.0	100.0
	Every day	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Methadone

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	65	.9	.9	.9
	Did not use	6878	98.7	98.7	99.6
	Once	8	.1	.1	99.8
	Every other month	6	.1	.1	99.8
	Once a month	4	.1	.1	99.9
	Once a week	5	.1	.1	100.0
	3 times a week	1	.0	.0	100.0
	Every day	1	.0	.0	100.0
	Total	6968	100.0	100.0	

Use of substance in the past 6 months without a prescription: Steroids

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	100	1.4	1.4	1.4
	Did not use	6852	98.3	98.3	99.8
	Once	7	.1	.1	99.9
	Every other month	1	.0	.0	99.9
	Once a month	1	.0	.0	99.9
	Once a week	1	.0	.0	99.9
	3 times a week	3	.0	.0	100.0
	5 times a week	1	.0	.0	100.0
	Every day	2	.0	.0	100.0
	Total	6968	100.0	100.0	

Alcohol use in the past 6 months

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	23	.3	.3	.3
	Yes	5651	81.1	81.1	81.4
	No	1294	18.6	18.6	100.0
	Total	6968	100.0	100.0	

Reasons for drinking: To experiment

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	18	.3	.3	.3
	Yes	1245	22.0	22.0	22.4
	No	4315	76.4	76.4	98.7
	Unsure	73	1.3	1.3	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: Because I am hooked

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.4	.4	.4
	Yes	132	2.3	2.3	2.8
	No	5415	95.8	95.8	98.6
	Unsure	80	1.4	1.4	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To relax or relieve tension

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Yes	3475	61.5	61.5	61.8
	No	2097	37.1	37.1	98.9
	Unsure	60	1.1	1.1	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To feel good or get high

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.4	.4	.4
	Yes	2831	50.1	50.1	50.5
	No	2728	48.3	48.3	98.8
	Unsure	68	1.2	1.2	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To seek deeper insights and understanding

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	42	.7	.7	.7
	Yes	353	6.2	6.2	7.0
	No	5173	91.5	91.5	98.5
	Unsure	83	1.5	1.5	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To have a good time with friends

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.1	.1	.1
	Yes	5016	88.8	88.8	88.9
	No	597	10.6	10.6	99.4
	Unsure	32	.6	.6	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To fit in with a group I like

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	18	.3	.3	.3
	Yes	818	14.5	14.5	14.8
	No	4748	84.0	84.0	98.8
	Unsure	67	1.2	1.2	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: Because of boredom

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Yes	1177	20.8	20.8	21.2
	No	4395	77.8	77.8	98.9
	Unsure	60	1.1	1.1	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: Because of anger or frustration

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Yes	850	15.0	15.0	15.4
	No	4740	83.9	83.9	99.3
	Unsure	42	.7	.7	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To get through the day

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.4	.4	.4
	Yes	206	3.6	3.6	4.1
	No	5388	95.3	95.3	99.4
	Unsure	33	.6	.6	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To increase the effects of other drug(s)

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	25	.4	.4	.4
	Yes	223	3.9	3.9	4.4
	No	5381	95.2	95.2	99.6
	Unsure	22	.4	.4	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To decrease the effects of other drug(s)

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	26	.5	.5	.5
	Yes	91	1.6	1.6	2.1
	No	5511	97.5	97.5	99.6
	Unsure	23	.4	.4	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To get to sleep

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.5	.5	.5
	Yes	550	9.7	9.7	10.2
	No	5042	89.2	89.2	99.4
	Unsure	32	.6	.6	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: Because it tastes good

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.4	.4	.4
	Yes	3145	55.7	55.7	56.1
	No	2420	42.8	42.8	98.9
	Unsure	62	1.1	1.1	100.0
	Total	5651	100.0	100.0	

Reasons for drinking: To get away from my problems

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	39	.7	.7	.7
	Yes	772	13.7	13.7	14.4
	No	4760	84.2	84.2	98.6
	Unsure	80	1.4	1.4	100.0
	Total	5651	100.0	100.0	

Combined alcohol with energy drinks

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.1	.1	.1
	Yes	1971	34.9	34.9	35.0
	No	3674	65.0	65.0	100.0
	Total	5651	100.0	100.0	

Combined alcohol with stimulants

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	3	.1	.1	.1
	Yes	386	6.8	6.8	6.9
	No	5262	93.1	93.1	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Friends or family members worried or complained about your drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	10	.2	.2	.2
	Never	4860	86.0	86.0	86.2
	Once	415	7.3	7.3	93.5
	Twice	182	3.2	3.2	96.7
	3-5 times	138	2.4	2.4	99.2
	6-9 times	19	.3	.3	99.5
	10+ times	27	.5	.5	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Had a hangover

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.5	.5	.5
	Never	1907	33.7	33.7	34.2
	Once	1082	19.1	19.1	53.4
	Twice	780	13.8	13.8	67.2
	3-5 times	953	16.9	16.9	84.0
	6-9 times	394	7.0	7.0	91.0
	10+ times	508	9.0	9.0	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Felt bad about your drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	40	.7	.7	.7
	Never	3842	68.0	68.0	68.7
	Once	836	14.8	14.8	83.5
	Twice	421	7.5	7.5	90.9
	3-5 times	350	6.2	6.2	97.1
	6-9 times	96	1.7	1.7	98.8
	10+ times	66	1.2	1.2	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Been unable to stop drinking when you wanted to

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Never	5189	91.8	91.8	92.2
	Once	152	2.7	2.7	94.9
	Twice	73	1.3	1.3	96.1
	3-5 times	103	1.8	1.8	98.0
	6-9 times	42	.7	.7	98.7
	10+ times	73	1.3	1.3	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Created problems between you and your friends or family members

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	23	.4	.4	.4
	Never	4839	85.6	85.6	86.0
	Once	407	7.2	7.2	93.2
	Twice	207	3.7	3.7	96.9
	3-5 times	141	2.5	2.5	99.4
	6-9 times	15	.3	.3	99.7
	10+ times	19	.3	.3	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Gotten into trouble at work or school because of drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	21	.4	.4	.4
	Never	5384	95.3	95.3	95.6
	Once	162	2.9	2.9	98.5
	Twice	54	1.0	1.0	99.5
	3-5 times	22	.4	.4	99.9
	6-9 times	1	.0	.0	99.9
	10+ times	7	.1	.1	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Gone to anyone for help to reduce your drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	18	.3	.3	.3
	Never	5518	97.6	97.6	98.0
	Once	72	1.3	1.3	99.2
	Twice	25	.4	.4	99.7
	3-5 times	11	.2	.2	99.9
	6-9 times	3	.1	.1	99.9
	10+ times	4	.1	.1	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Been hurt or injured because of drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	13	.2	.2	.2
	Never	4899	86.7	86.7	86.9
	Once	424	7.5	7.5	94.4
	Twice	169	3.0	3.0	97.4
	3-5 times	103	1.8	1.8	99.2
	6-9 times	29	.5	.5	99.8
	10+ times	14	.2	.2	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Been ticketed or arrested because of drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	19	.3	.3	.3
	Never	5444	96.3	96.3	96.7
	Once	160	2.8	2.8	99.5
	Twice	19	.3	.3	99.8
	3-5 times	4	.1	.1	99.9
	6-9 times	2	.0	.0	99.9
	10+ times	3	.1	.1	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Been arrested for drunk driving

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	21	.4	.4	.4
	Never	5601	99.1	99.1	99.5
	Once	20	.4	.4	99.8
	Twice	6	.1	.1	99.9
	3-5 times	2	.0	.0	100.0
	10+ times	1	.0	.0	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Driven a car while under the influence

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.5	.5	.5
	Never	4861	86.0	86.0	86.5
	Once	337	6.0	6.0	92.5
	Twice	212	3.8	3.8	96.2
	3-5 times	143	2.5	2.5	98.7
	6-9 times	28	.5	.5	99.2
	10+ times	43	.8	.8	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Gotten into physical fights when drinking

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	52	.9	.9	.9
	Never	5337	94.4	94.4	95.4
	Once	171	3.0	3.0	98.4
	Twice	61	1.1	1.1	99.5
	3-5 times	22	.4	.4	99.9
	6-9 times	4	.1	.1	99.9
	10+ times	4	.1	.1	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Experienced unwanted sexual attention

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	33	.6	.6	.6
	Never	4990	88.3	88.3	88.9
	Once	346	6.1	6.1	95.0
	Twice	148	2.6	2.6	97.6
	3-5 times	96	1.7	1.7	99.3
	6-9 times	19	.3	.3	99.7
	10+ times	19	.3	.3	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Engaged in risky sexual behavior

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	18	.3	.3	.3
	Never	4856	85.9	85.9	86.3
	Once	357	6.3	6.3	92.6
	Twice	205	3.6	3.6	96.2
	3-5 times	149	2.6	2.6	98.8
	6-9 times	33	.6	.6	99.4
	10+ times	33	.6	.6	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Missed class or assignments

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	25	.4	.4	.4
	Never	4500	79.6	79.6	80.1
	Once	452	8.0	8.0	88.1
	Twice	318	5.6	5.6	93.7
	3-5 times	247	4.4	4.4	98.1
	6-9 times	60	1.1	1.1	99.1
	10+ times	49	.9	.9	100.0
	Total	5651	100.0	100.0	

Consequences of alcohol use: Blacked out

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	26	.5	.5	.5
	Never	3734	66.1	66.1	66.5
	Once	762	13.5	13.5	80.0
	Twice	439	7.8	7.8	87.8
	3-5 times	379	6.7	6.7	94.5
	6-9 times	164	2.9	2.9	97.4
	10+ times	147	2.6	2.6	100.0
	Total	5651	100.0	100.0	

Usual type of alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	67	1.2	1.2	1.2
	Beer	1452	25.7	25.7	26.9
	Malt beverages	347	6.1	6.1	33.0
	Wine	646	11.4	11.4	44.5
	Liquor	3012	53.3	53.3	97.8
	High alcohol drinks	54	1.0	1.0	98.7
	Some other type	73	1.3	1.3	100.0
	Total	5651	100.0	100.0	

Age

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	21 or older	3213	46.1	46.1	46.1
	Under 21	3755	53.9	53.9	100.0
	Total	6968	100.0	100.0	

Students under 21: 1st easiest way to get alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	90	3.2	3.2	3.2
	Older friends	1476	52.2	52.2	55.4
	Older siblings	91	3.2	3.2	58.6
	On-campus parties	121	4.3	4.3	62.9
	Fraternity or sorority houses	438	15.5	15.5	78.4
	Off-campus private parties	301	10.6	10.6	89.0
	Someone else on campus	128	4.5	4.5	93.6
	Buying it yourself	46	1.6	1.6	95.2
	Stealing it from store	4	.1	.1	95.3
	From parents without their knowledge	13	.5	.5	95.8
	From parents with their knowledge	119	4.2	4.2	100.0
	Total	2827	100.0	100.0	

Students under 21: 2nd easiest way to get alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	161	5.7	5.7	5.7
	Older friends	505	17.9	17.9	23.6
	Older siblings	271	9.6	9.6	33.1
	On-campus parties	332	11.7	11.7	44.9
	Fraternity or sorority houses	331	11.7	11.7	56.6
	Off-campus private parties	763	27.0	27.0	83.6
	Someone else on campus	245	8.7	8.7	92.3
	Buying it yourself	52	1.8	1.8	94.1
	Stealing it from store	3	.1	.1	94.2
	From parents without their knowledge	31	1.1	1.1	95.3
	From parents with their knowledge	133	4.7	4.7	100.0
	Total	2827	100.0	100.0	

Students under 21: 3rd easiest way to get alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	238	8.4	8.4	8.4
	Older friends	328	11.6	11.6	20.0
	Older siblings	212	7.5	7.5	27.5
	On-campus parties	362	12.8	12.8	40.3
	Fraternity or sorority houses	322	11.4	11.4	51.7
	Off-campus private parties	509	18.0	18.0	69.7
	Someone else on campus	452	16.0	16.0	85.7
	Buying it yourself	82	2.9	2.9	88.6
	Stealing it from store	11	.4	.4	89.0
	From parents without their knowledge	78	2.8	2.8	91.8
	From parents with their knowledge	233	8.2	8.2	100.0
	Total	2827	100.0	100.0	

Students under 21: Purchased alcohol from retailer

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	20	.7	.7	.7
	Yes	287	10.2	10.2	10.9
	No	2520	89.1	89.1	100.0
	Total	2827	100.0	100.0	

Students under 21: Most common method to purchase alcohol from retailer

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	2.1	2.1	2.1
	Used a fake ID	90	31.4	31.4	33.4
	Used someone elses ID	71	24.7	24.7	58.2
	Bought alcohol without using an ID	120	41.8	41.8	100.0
	Total	287	100.0	100.0	

Students under 21: Easiest retail place to purchase alcohol

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	2	.7	.7	.7
	Bar	81	28.2	28.2	28.9
	Restaurant	75	26.1	26.1	55.1
	Grocery store	33	11.5	11.5	66.6
	Discount store	18	6.3	6.3	72.8
	Convenience store	24	8.4	8.4	81.2
	Drug store	13	4.5	4.5	85.7
	Liquor store	39	13.6	13.6	99.3
	Winery or microbrewery	2	.7	.7	100.0
	Total	287	100.0	100.0	

Any drug use in the past 6 months

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	5	.1	.1	.1
	Yes	1523	21.9	21.9	21.9
	No	5440	78.1	78.1	100.0
	Total	6968	100.0	100.0	

Consequences of drug use: Abused more than one drug at a time

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	8	.5	.5	.5
	Never	993	65.2	65.2	65.7
	Once	114	7.5	7.5	73.2
	Twice	91	6.0	6.0	79.2
	3-5 times	153	10.0	10.0	89.2
	6-9 times	51	3.3	3.3	92.6
	10+ times	113	7.4	7.4	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Been unable to stop using drugs when you wanted to

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.4	.4	.4
	Never	1385	90.9	90.9	91.3
	Once	33	2.2	2.2	93.5
	Twice	21	1.4	1.4	94.9
	3-5 times	27	1.8	1.8	96.7
	6-9 times	12	.8	.8	97.4
	10+ times	39	2.6	2.6	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Had blackouts or flashbacks as a result of drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.4	.4	.4
	Never	1317	86.5	86.5	86.9
	Once	69	4.5	4.5	91.4
	Twice	41	2.7	2.7	94.1
	3-5 times	45	3.0	3.0	97.0
	6-9 times	23	1.5	1.5	98.6
	10+ times	22	1.4	1.4	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Felt bad or guilty about your drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.4	.4	.4
	Never	1072	70.4	70.4	70.8
	Once	159	10.4	10.4	81.2
	Twice	114	7.5	7.5	88.7
	3-5 times	107	7.0	7.0	95.7
	6-9 times	22	1.4	1.4	97.2
	10+ times	43	2.8	2.8	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Had your friends or family complain about your drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.4	.4	.4
	Never	1328	87.2	87.2	87.6
	Once	79	5.2	5.2	92.8
	Twice	49	3.2	3.2	96.0
	3-5 times	36	2.4	2.4	98.4
	6-9 times	9	.6	.6	98.9
	10+ times	16	1.1	1.1	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Created problems between you and your friends or family members

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	7	.5	.5	.5
	Never	1379	90.5	90.5	91.0
	Once	67	4.4	4.4	95.4
	Twice	34	2.2	2.2	97.6
	3-5 times	21	1.4	1.4	99.0
	6-9 times	4	.3	.3	99.3
	10+ times	11	.7	.7	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Engaged in illegal activities in order to obtain drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	7	.5	.5	.5
	Never	1333	87.5	87.5	88.0
	Once	34	2.2	2.2	90.2
	Twice	25	1.6	1.6	91.9
	3-5 times	37	2.4	2.4	94.3
	6-9 times	19	1.2	1.2	95.5
	10+ times	68	4.5	4.5	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Gotten into physical fights because you were using drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	4	.3	.3	.3
	Never	1493	98.0	98.0	98.3
	Once	10	.7	.7	98.9
	Twice	6	.4	.4	99.3
	3-5 times	4	.3	.3	99.6
	6-9 times	1	.1	.1	99.7
	10+ times	5	.3	.3	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Experienced withdrawal symptoms when you stopped taking drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	8	.5	.5	.5
	Never	1394	91.5	91.5	92.1
	Once	32	2.1	2.1	94.2
	Twice	37	2.4	2.4	96.6
	3-5 times	29	1.9	1.9	98.5
	6-9 times	8	.5	.5	99.0
	10+ times	15	1.0	1.0	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Had medical problems as a result of your drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	5	.3	.3	.3
	Never	1436	94.3	94.3	94.6
	Once	32	2.1	2.1	96.7
	Twice	20	1.3	1.3	98.0
	3-5 times	12	.8	.8	98.8
	6-9 times	7	.5	.5	99.3
	10+ times	11	.7	.7	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Driven a car while under the influence of drugs

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	7	.5	.5	.5
	Never	955	62.7	62.7	63.2
	Once	110	7.2	7.2	70.4
	Twice	103	6.8	6.8	77.2
	3-5 times	121	7.9	7.9	85.1
	6-9 times	42	2.8	2.8	87.9
	10+ times	185	12.1	12.1	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Gotten in trouble at work or school because of drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	11	.7	.7	.7
	Never	1465	96.2	96.2	96.9
	Once	26	1.7	1.7	98.6
	Twice	6	.4	.4	99.0
	3-5 times	6	.4	.4	99.4
	6-9 times	4	.3	.3	99.7
	10+ times	5	.3	.3	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Been in very vulnerable situations

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	7	.5	.5	.5
	Never	1420	93.2	93.2	93.7
	Once	41	2.7	2.7	96.4
	Twice	21	1.4	1.4	97.8
	3-5 times	17	1.1	1.1	98.9
	6-9 times	4	.3	.3	99.1
	10+ times	13	.9	.9	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Engaged in risky sexual behavior

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	8	.5	.5	.5
	Never	1384	90.9	90.9	91.4
	Once	46	3.0	3.0	94.4
	Twice	29	1.9	1.9	96.3
	3-5 times	31	2.0	2.0	98.4
	6-9 times	6	.4	.4	98.8
	10+ times	19	1.2	1.2	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Failed a class because of drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	9	.6	.6	.6
	Never	1489	97.8	97.8	98.4
	Once	15	1.0	1.0	99.3
	Twice	5	.3	.3	99.7
	3-5 times	3	.2	.2	99.9
	6-9 times	1	.1	.1	99.9
	10+ times	1	.1	.1	100.0
	Total	1523	100.0	100.0	

Consequences of drug use: Missed class or assignments due to drug use

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6	.4	.4	.4
	Never	1266	83.1	83.1	83.5
	Once	84	5.5	5.5	89.0
	Twice	56	3.7	3.7	92.7
	3-5 times	69	4.5	4.5	97.2
	6-9 times	20	1.3	1.3	98.6
	10+ times	22	1.4	1.4	100.0
	Total	1523	100.0	100.0	

Perception of peer approval/disapproval: Trying marijuana once or twice

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	28	.4	.4	.4
	Strongly approve	1573	22.6	22.6	23.0
	Somewhat approve	2650	38.0	38.0	61.0
	Neither approve or disapprove	1763	25.3	25.3	86.3
	Somewhat disapprove	461	6.6	6.6	92.9
	Strongly disapprove	493	7.1	7.1	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Smoking marijuana occasionally

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	42	.6	.6	.6
	Strongly approve	954	13.7	13.7	14.3
	Somewhat approve	2454	35.2	35.2	49.5
	Neither approve or disapprove	2065	29.6	29.6	79.1
	Somewhat disapprove	826	11.9	11.9	91.0
	Strongly disapprove	627	9.0	9.0	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Smoking marijuana regularly

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	45	.6	.6	.6
	Strongly approve	412	5.9	5.9	6.6
	Somewhat approve	1330	19.1	19.1	25.6
	Neither approve or disapprove	2018	29.0	29.0	54.6
	Somewhat disapprove	1969	28.3	28.3	82.9
	Strongly disapprove	1194	17.1	17.1	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Trying cocaine once or twice

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	40	.6	.6	.6
	Strongly approve	110	1.6	1.6	2.2
	Somewhat approve	564	8.1	8.1	10.2
	Neither approve or disapprove	1118	16.0	16.0	26.3
	Somewhat disapprove	2497	35.8	35.8	62.1
	Strongly disapprove	2639	37.9	37.9	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Taking cocaine regularly

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	59	.8	.8	.8
	Strongly approve	63	.9	.9	1.8
	Somewhat approve	135	1.9	1.9	3.7
	Neither approve or disapprove	639	9.2	9.2	12.9
	Somewhat disapprove	1269	18.2	18.2	31.1
	Strongly disapprove	4803	68.9	68.9	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Trying amphetamines once or twice

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	37	.5	.5	.5
	Strongly approve	715	10.3	10.3	10.8
	Somewhat approve	1661	23.8	23.8	34.6
	Neither approve or disapprove	1901	27.3	27.3	61.9
	Somewhat disapprove	1173	16.8	16.8	78.7
	Strongly disapprove	1481	21.3	21.3	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Taking amphetamines regularly

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	73	1.0	1.0	1.0
	Strongly approve	272	3.9	3.9	5.0
	Somewhat approve	749	10.7	10.7	15.7
	Neither approve or disapprove	1515	21.7	21.7	37.4
	Somewhat disapprove	1676	24.1	24.1	61.5
	Strongly disapprove	2683	38.5	38.5	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Having one or two alcoholic drinks nearly every day

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	42	.6	.6	.6
	Strongly approve	732	10.5	10.5	11.1
	Somewhat approve	1573	22.6	22.6	33.7
	Neither approve or disapprove	2122	30.5	30.5	64.1
	Somewhat disapprove	1678	24.1	24.1	88.2
	Strongly disapprove	821	11.8	11.8	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Having three or four alcoholic drinks nearly every day

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	42	.6	.6	.6
	Strongly approve	316	4.5	4.5	5.1
	Somewhat approve	808	11.6	11.6	16.7
	Neither approve or disapprove	1534	22.0	22.0	38.7
	Somewhat disapprove	2140	30.7	30.7	69.5
	Strongly disapprove	2128	30.5	30.5	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Having five or more drinks in one sitting

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	52	.7	.7	.7
	Strongly approve	851	12.2	12.2	13.0
	Somewhat approve	1466	21.0	21.0	34.0
	Neither approve or disapprove	1911	27.4	27.4	61.4
	Somewhat disapprove	1247	17.9	17.9	79.3
	Strongly disapprove	1441	20.7	20.7	100.0
	Total	6968	100.0	100.0	

Perception of peer approval/disapproval: Taking steroids for body building or improved athletic performance

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	51	.7	.7	.7
	Strongly approve	108	1.5	1.5	2.3
	Somewhat approve	423	6.1	6.1	8.4
	Neither approve or disapprove	1622	23.3	23.3	31.6
	Somewhat disapprove	2047	29.4	29.4	61.0
	Strongly disapprove	2717	39.0	39.0	100.0
	Total	6968	100.0	100.0	

Perceived number of alcoholic drinks consumed by the typical student at the last party

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	36	.5	.5	.5
	0	157	2.3	2.3	2.8
	1	82	1.2	1.2	3.9
	2	312	4.5	4.5	8.4
	3	810	11.6	11.6	20.0
	4	954	13.7	13.7	33.7
	5	1383	19.8	19.8	53.6
	6	937	13.4	13.4	67.0
	7	654	9.4	9.4	76.4
	8	549	7.9	7.9	84.3
	9	160	2.3	2.3	86.6
	10	596	8.6	8.6	95.1
	11	25	.4	.4	95.5
	12	130	1.9	1.9	97.4
	13	27	.4	.4	97.8
	14	24	.3	.3	98.1
	15	61	.9	.9	99.0
	16	14	.2	.2	99.2
	17	5	.1	.1	99.3
	18	9	.1	.1	99.4
	19	4	.1	.1	99.4
	20	16	.2	.2	99.7
	21	2	.0	.0	99.7
	22	2	.0	.0	99.7
	23	2	.0	.0	99.8
	24	1	.0	.0	99.8
	25	2	.0	.0	99.8
	27	1	.0	.0	99.8
	28	2	.0	.0	99.8
	30	11	.2	.2	100.0
	Total	6968	100.0	100.0	

Gambling: Played cards at home

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	30	.4	.4	.4
	Never	5877	84.3	84.3	84.8
	1-2 times per year	579	8.3	8.3	93.1
	3-6 times per year	329	4.7	4.7	97.8
	Monthly	109	1.6	1.6	99.4
	Weekly	35	.5	.5	99.9
	3+ times per week	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Gambling: Gambled at a casino

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	32	.5	.5	.5
	Never	6023	86.4	86.4	86.9
	1-2 times per year	709	10.2	10.2	97.1
	3-6 times per year	174	2.5	2.5	99.6
	Monthly	18	.3	.3	99.8
	Weekly	8	.1	.1	99.9
	3+ times per week	4	.1	.1	100.0
	Total	6968	100.0	100.0	

Gambling: Gambled on-line

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	44	.6	.6	.6
	Never	6783	97.3	97.3	98.0
	1-2 times per year	68	1.0	1.0	99.0
	3-6 times per year	28	.4	.4	99.4
	Monthly	17	.2	.2	99.6
	Weekly	14	.2	.2	99.8
	3+ times per week	14	.2	.2	100.0
	Total	6968	100.0	100.0	

Gambling: Gambled at informal locations

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	38	.5	.5	.5
	Never	6836	98.1	98.1	98.7
	1-2 times per year	48	.7	.7	99.3
	3-6 times per year	22	.3	.3	99.7
	Monthly	15	.2	.2	99.9
	Weekly	4	.1	.1	99.9
	3+ times per week	5	.1	.1	100.0
	Total	6968	100.0	100.0	

Gambling: Bet on fantasy football/basketball

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	27	.4	.4	.4
	Never	6358	91.2	91.2	91.6
	1-2 times per year	405	5.8	5.8	97.4
	3-6 times per year	99	1.4	1.4	98.9
	Monthly	41	.6	.6	99.5
	Weekly	29	.4	.4	99.9
	3+ times per week	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Gambling: Betting pools

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	41	.6	.6	.6
	Never	6097	87.5	87.5	88.1
	1-2 times per year	689	9.9	9.9	98.0
	3-6 times per year	95	1.4	1.4	99.3
	Monthly	28	.4	.4	99.7
	Weekly	11	.2	.2	99.9
	3+ times per week	7	.1	.1	100.0
	Total	6968	100.0	100.0	

Gambling: Lottery

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	35	.5	.5	.5
	Never	5218	74.9	74.9	75.4
	1-2 times per year	1052	15.1	15.1	90.5
	3-6 times per year	490	7.0	7.0	97.5
	Monthly	125	1.8	1.8	99.3
	Weekly	37	.5	.5	99.8
	3+ times per week	11	.2	.2	100.0
	Total	6968	100.0	100.0	

Gambling: Placed bet with bookie

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	63	.9	.9	.9
	Never	6799	97.6	97.6	98.5
	1-2 times per year	44	.6	.6	99.1
	3-6 times per year	24	.3	.3	99.5
	Monthly	15	.2	.2	99.7
	Weekly	9	.1	.1	99.8
	3+ times per week	14	.2	.2	100.0
	Total	6968	100.0	100.0	

Location where you gambled most frequently

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	160	2.3	2.3	2.3
	I did not gamble in the past year	4843	69.5	69.5	71.8
	I only gambled online or playing the lottery	527	7.6	7.6	79.4
	Apartment/House	484	6.9	6.9	86.3
	Residence hall	55	.8	.8	87.1
	Fraternity/Sorority	42	.6	.6	87.7
	Other campus housing	8	.1	.1	87.8
	Casino	649	9.3	9.3	97.1
	Other	200	2.9	2.9	100.0
	Total	6968	100.0	100.0	

Consequences of gambling: Academic problems

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	20	.7	.7	.7
	I do not gamble	1165	41.4	41.4	42.2
	None	1619	57.6	57.6	99.8
	Occasionally	4	.1	.1	99.9
	Frequently	3	.1	.1	100.0
	Total	2811	100.0	100.0	

Consequences of gambling: Issues with legal system

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	20	.7	.7	.7
	I do not gamble	1163	41.4	41.4	42.1
	None	1623	57.7	57.7	99.8
	Occasionally	2	.1	.1	99.9
	Frequently	3	.1	.1	100.0
	Total	2811	100.0	100.0	

Consequences of gambling: Financial issues

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.9	.9	.9
	I do not gamble	1157	41.2	41.2	42.0
	None	1582	56.3	56.3	98.3
	Occasionally	45	1.6	1.6	99.9
	Frequently	3	.1	.1	100.0
	Total	2811	100.0	100.0	

Consequences of gambling: Felt bad about gambling

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	29	1.0	1.0	1.0
	I do not gamble	1155	41.1	41.1	42.1
	None	1521	54.1	54.1	96.2
	Occasionally	99	3.5	3.5	99.8
	Frequently	7	.2	.2	100.0
	Total	2811	100.0	100.0	

Consequences of gambling: Poor physical health

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	24	.9	.9	.9
	I do not gamble	1161	41.3	41.3	42.2
	None	1616	57.5	57.5	99.6
	Occasionally	5	.2	.2	99.8
	Frequently	5	.2	.2	100.0
	Total	2811	100.0	100.0	

Consequences of gambling: Issues with friends and family

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	32	1.1	1.1	1.1
	I do not gamble	1159	41.2	41.2	42.4
	None	1594	56.7	56.7	99.1
	Occasionally	20	.7	.7	99.8
	Frequently	6	.2	.2	100.0
	Total	2811	100.0	100.0	

Number of days in past month that mental health was not good

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	98	1.4	1.4	1.4
	0	1633	23.4	23.4	24.8
	1	461	6.6	6.6	31.5
	2	734	10.5	10.5	42.0
	3	598	8.6	8.6	50.6
	4	384	5.5	5.5	56.1
	5	586	8.4	8.4	64.5
	6	140	2.0	2.0	66.5
	7	228	3.3	3.3	69.8
	8	112	1.6	1.6	71.4
	9	57	.8	.8	72.2
	10	399	5.7	5.7	77.9
	11	20	.3	.3	78.2
	12	68	1.0	1.0	79.2
	13	28	.4	.4	79.6
	14	35	.5	.5	80.1
	15	216	3.1	3.1	83.2

Number of days in past month that mental health was not good

	Frequency	Percent	Valid Percent	Cumulative Percent
16	17	.2	.2	83.4
17	30	.4	.4	83.9
18	15	.2	.2	84.1
19	5	.1	.1	84.2
20	151	2.2	2.2	86.3
21	17	.2	.2	86.6
22	14	.2	.2	86.8
23	10	.1	.1	86.9
24	21	.3	.3	87.2
25	84	1.2	1.2	88.4
26	14	.2	.2	88.6
27	14	.2	.2	88.8
28	29	.4	.4	89.2
29	13	.2	.2	89.4
30	119	1.7	1.7	91.1
Dont Know	618	8.9	8.9	100.0
Total	6968	100.0	100.0	

Number of days in past 2 weeks that you had suicidal thoughts or thoughts of hurting yourself

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	322	4.6	4.6	4.6
	0	5745	82.4	82.4	87.1
	1	339	4.9	4.9	91.9
	2	205	2.9	2.9	94.9
	3	96	1.4	1.4	96.3
	4	55	.8	.8	97.0
	5	42	.6	.6	97.6
	6	24	.3	.3	98.0
	7	41	.6	.6	98.6
	8	9	.1	.1	98.7
	9	7	.1	.1	98.8
	10	30	.4	.4	99.2
	11	5	.1	.1	99.3
	12	12	.2	.2	99.5
	13	8	.1	.1	99.6
	14	28	.4	.4	100.0
Total		6968	100.0	100.0	

Gender

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Male	2572	36.9	36.9	36.9
	Female	4396	63.1	63.1	100.0
Total		6968	100.0	100.0	

Sexual identification

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	17	.2	.2	.2
	Heterosexual	6409	92.0	92.0	92.2
	Gay or Lesbian	175	2.5	2.5	94.7
	Bi-sexual	220	3.2	3.2	97.9
	Other	48	.7	.7	98.6
	Do not know/Not sure	99	1.4	1.4	100.0
Total		6968	100.0	100.0	

Ethnicity

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	55	.8	.8	.8
	Hispanic	259	3.7	3.7	4.5
	Non-Hispanic	6654	95.5	95.5	100.0
	Total	6968	100.0	100.0	

Race

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	45	.6	.6	.6
	White	5879	84.4	84.4	85.0
	Black	239	3.4	3.4	88.4
	Asian	502	7.2	7.2	95.7
	Native American or Alaskan Native	28	.4	.4	96.1
	Hawaiian or Pacific Islander	9	.1	.1	96.2
	More than one race	163	2.3	2.3	98.5
	Other	103	1.5	1.5	100.0
	Total	6968	100.0	100.0	

Marital status

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	25	.4	.4	.4
	Single, never married	6436	92.4	92.4	92.7
	Single, living with partner	291	4.2	4.2	96.9
	Married	168	2.4	2.4	99.3
	Separated	7	.1	.1	99.4
	Divorced	32	.5	.5	99.9
	Widowed	9	.1	.1	100.0
	Total	6968	100.0	100.0	

Student status

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	41	.6	.6	.6
	Full-time	6758	97.0	97.0	97.6
	Part-time	169	2.4	2.4	100.0
	Total	6968	100.0	100.0	

Classification

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	21	.3	.3	.3
	Freshman or first year	1794	25.7	25.7	26.0
	Sophomore or second year	1677	24.1	24.1	50.1
	Junior	1655	23.8	23.8	73.9
	Senior	1645	23.6	23.6	97.5
	Grad or professional	108	1.5	1.5	99.0
	Not seeking a degree	27	.4	.4	99.4
	Other	41	.6	.6	100.0
	Total	6968	100.0	100.0	

Location of current residence

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	46	.7	.7	.7
	On-campus	3816	54.8	54.8	55.4
	Off-campus	3106	44.6	44.6	100.0
	Total	6968	100.0	100.0	

Housing type

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	37	.5	.5	.5
	House or apartment	3393	48.7	48.7	49.2
	Residence hall	2707	38.8	38.8	88.1
	Fraternity or sorority	530	7.6	7.6	95.7
	Other campus housing	176	2.5	2.5	98.2
	Other	125	1.8	1.8	100.0
	Total	6968	100.0	100.0	

With whom do you live: Alone

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	5966	85.6	85.6	85.6
	Yes	1002	14.4	14.4	100.0
	Total	6968	100.0	100.0	

With whom do you live: With roomates

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	1617	23.2	23.2	23.2
	Yes	5351	76.8	76.8	100.0
	Total	6968	100.0	100.0	

With whom do you live: With parents

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6607	94.8	94.8	94.8
	Yes	361	5.2	5.2	100.0
	Total	6968	100.0	100.0	

With whom do you live: With spouse

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6746	96.8	96.8	96.8
	Yes	222	3.2	3.2	100.0
	Total	6968	100.0	100.0	

With whom do you live: With children

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6846	98.2	98.2	98.2
	Yes	122	1.8	1.8	100.0
	Total	6968	100.0	100.0	

With whom do you live: Other

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	6812	97.8	97.8	97.8
	Yes	156	2.2	2.2	100.0
	Total	6968	100.0	100.0	

Employment status

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	28	.4	.4	.4
	Full-time	249	3.6	3.6	4.0
	Part-time	3043	43.7	43.7	47.6
	None	3648	52.4	52.4	100.0
	Total	6968	100.0	100.0	

Military status

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	No answer	21	.3	.3	.3
	Have served	62	.9	.9	1.2
	Have not served	6885	98.8	98.8	100.0
	Total	6968	100.0	100.0	

Indiana College Substance Use Survey

2012

1. How many times in the past **year** have you used...? (Mark one for each line.)

	Never	1-5 times	6-19 times	20-39 times	40 times or more
a. Cigarettes	A	B	C	D	E
b. Cigars	A	B	C	D	E
c. Chewing/smokeless tobacco	A	B	C	D	E
d. Smoking tobacco with hookah/water pipe	A	B	C	D	E
e. Alcohol (beer, wine, liquor)	A	B	C	D	E
f. Marijuana (pot, hashish, hash oil)	A	B	C	D	E
g. Cocaine (crack, rock, freebase)	A	B	C	D	E
h. Hallucinogens (LSD, PCP, ecstasy)	A	B	C	D	E
i. Heroin	A	B	C	D	E
j. Inhalants (glue, solvents, gas)	A	B	C	D	E
k. Prescription pills that are not prescribed to you	A	B	C	D	E
l. Medications prescribed to you, but you used to get high	A	B	C	D	E
m. Over-the-counter medications to get high	A	B	C	D	E
n. Methamphetamine	A	B	C	D	E
o. Other illegal drugs	A	B	C	D	E

2. How many times in the past **month** have you used...? (Mark one for each line.)

	Never	1-5 times	6-19 times	20-39 times	40 times or more
a. Cigarettes	A	B	C	D	E
b. Cigars	A	B	C	D	E
c. Chewing/smokeless tobacco	A	B	C	D	E
d. Smoking tobacco with hookah/water pipe	A	B	C	D	E
e. Alcohol (beer, wine, liquor)	A	B	C	D	E
f. Marijuana (pot, hashish, hash oil)	A	B	C	D	E
g. Cocaine (crack, rock, freebase)	A	B	C	D	E
h. Hallucinogens (LSD, PCP, ecstasy)	A	B	C	D	E
i. Heroin	A	B	C	D	E
j. Inhalants (glue, solvents, gas)	A	B	C	D	E
k. Prescription pills that are not prescribed to you	A	B	C	D	E
l. Medications prescribed to you, but you used to get high	A	B	C	D	E
m. Over-the-counter medications to get high	A	B	C	D	E
n. Methamphetamine	A	B	C	D	E
o. Other illegal drugs	A	B	C	D	E

**Use this definition for the following two questions.
One drink is a 12-ounce bottle of beer, a 5-ounce glass of wine, or
one shot of liquor straight or in a mixed drink.**

3. During the **past 30 days**, on how many **days** did you drink 5 or more alcoholic drinks (for males) or 4 or more alcoholic drinks (for females) in 2-3 hours or less? (If zero, enter 00. If less than 10, enter 01, 02, 03, etc.)

--	--

4. Think back over the **last two weeks**. How many **times** have you had 5 or more alcoholic drinks (for males) or 4 or more alcoholic drinks (for females) in 2-3 hours or less? (If zero, enter 00. If less than 10, enter 01, 02, 03, etc.)

--	--

5. How old were you when you **first used**... (Mark one for each line.)

	Did not use	Under 10	10-11	12-13	14-15	16-17	18-20	21-25	26+
a. Cigarettes	A	B	C	D	E	F	G	H	I
b. Cigars	A	B	C	D	E	F	G	H	I
c. Chewing/smokeless tobacco	A	B	C	D	E	F	G	H	I
d. Smoked tobacco with hookah/water pipe	A	B	C	D	E	F	G	H	I
e. Alcohol (beer, wine, liquor)	A	B	C	D	E	F	G	H	I
f. Marijuana (pot, hashish, hash oil)	A	B	C	D	E	F	G	H	I
g. Cocaine (crack, rock, freebase)	A	B	C	D	E	F	G	H	I
h. Hallucinogens (LSD, PCP, ecstasy)	A	B	C	D	E	F	G	H	I
i. Heroin	A	B	C	D	E	F	G	H	I
j. Inhalants (glue, solvents, gas)	A	B	C	D	E	F	G	H	I
k. Prescription pills that are not prescribed to you	A	B	C	D	E	F	G	H	I
l. Medications prescribed to you, but you used to get high	A	B	C	D	E	F	G	H	I
m. Over-the-counter medications to get high	A	B	C	D	E	F	G	H	I
n. Methamphetamine	A	B	C	D	E	F	G	H	I
o. Other illegal drugs	A	B	C	D	E	F	G	H	I

6. **In the last 6 months**, how many times have you used the following drugs *without* a prescription? (Mark one for each line.)

	Did not use	Once	Every other month	Once/month	Once/week	3 times/week	5 times/week	Every day
a. Adderall	A	B	C	D	E	F	G	H
b. Ritalin	A	B	C	D	E	F	G	H
c. Xanax	A	B	C	D	E	F	G	H
d. Codeine	A	B	C	D	E	F	G	H
e. Vicodin	A	B	C	D	E	F	G	H
f. Lortab	A	B	C	D	E	F	G	H
g. Percocet	A	B	C	D	E	F	G	H
h. Oxycontin	A	B	C	D	E	F	G	H
i. Morphine	A	B	C	D	E	F	G	H
j. Methadone	A	B	C	D	E	F	G	H
k. Steroids	A	B	C	D	E	F	G	H

7. Did you drink alcohol in the **last 6 months**?

- A. Yes (Go to question #8)
 B. No (Go to Question #13)

8. Did you drink alcoholic beverages for any of the following reasons in the **last 6 months**? (Mark one for each line.)

	Yes	No	Unsure
a. To experiment (to see what it's like)	Y	N	U
b. Because I am "hooked" (I feel I have to drink)	Y	N	U
c. To relax or relieve tension	Y	N	U
d. To feel good or get high	Y	N	U
e. To seek deeper insights and understanding	Y	N	U
f. To have a good time with my friends	Y	N	U
g. To fit in with a group I like	Y	N	U
h. Because of boredom, nothing else to do	Y	N	U
i. Because of anger or frustration	Y	N	U
j. To get through the day	Y	N	U
k. To increase the effects of some other drug(s)	Y	N	U
l. To decrease (offset) the effects of some other drug(s)	Y	N	U
m. To get to sleep	Y	N	U
n. Because it tastes good	Y	N	U
o. To get away from my problems or troubles	Y	N	U

9. Have you combined alcohol with energy drinks (e.g., Red Bull, Monster) in the past 6 months?
 Y N

10. Have you combined alcohol with stimulants (e.g., Adderall, Ritalin) in the past 6 months?
 Y N

11. Have you experienced the following due to your **drinking in the last 6 months.** (Mark one for each line.)

	Number of Times					
	Never	Once	Twice	3-5 times	6-9 times	10+ times
a. Had friends or family members worry or complain about your drinking	A	B	C	D	E	F
b. Had a hangover	A	B	C	D	E	F
c. Felt bad or guilty about your drinking	A	B	C	D	E	F
d. Been unable to stop drinking when you wanted to	A	B	C	D	E	F
e. Created problems between you and your friends or family members	A	B	C	D	E	F
f. Gotten into trouble at work or school because of drinking	A	B	C	D	E	F
g. Gone to anyone for help to reduce your drinking	A	B	C	D	E	F
h. Been hurt or injured because of drinking	A	B	C	D	E	F
i. Been ticketed or arrested because of your drinking	A	B	C	D	E	F
j. Been arrested for drunk driving	A	B	C	D	E	F
k. Driven a car while under the influence	A	B	C	D	E	F
l. Gotten into physical fights when drinking	A	B	C	D	E	F
m. Experienced unwanted sexual attention such as being touched when you don't want to be, forced kissing, etc.	A	B	C	D	E	F
n. Engaged in risky sexual behavior such as having unprotected sex or sex with someone you just met because you were drinking	A	B	C	D	E	F
o. Missed class or an assignment	A	B	C	D	E	F
p. Forgot where you were or what you did (blacked out)	A	B	C	D	E	F

12. If you drank alcohol in the **past 6 months**, what type of alcohol did you **usually** drink? (Mark one only)

- a. Beer
- b. Malt beverages, such as Smirnoff Ice, Bacardi Silver, Hard Lemonade
- c. Wine
- d. Liquor, such as vodka, rum, whiskey
- e. High alcohol drinks, such as Everclear, Bacardi 151, 99 bananas
- f. Some other type

13. Are you 21 years or older?

- a. No, I am under 21 years old
- b. Yes

***** IF YOU ARE 21 OR OLDER, SKIP TO QUESTION #18**

*****IF YOU ARE UNDER 21 AND DID NOT DRINK ALCOHOL IN THE PAST YEAR, SKIP TO QUESTION #18*****

14. IF YOU ARE UNDER 21 and drink alcohol, what are the **three easiest ways** for you to get alcohol?

- A. older friends
- B. older siblings
- C. on-campus parties
- D. fraternity or sorority houses
- E. off-campus private parties (apartments, houses)
- F. someone else on-campus gives it to you
- G. buying it yourself
- H. stealing it from store
- I. from parents without their knowledge
- J. from parents with their knowledge

First easiest: _____ **Second easiest:** _____ **Third easiest:** _____

(Please place your choice on the response sheet)

15. IF YOU ARE UNDER 21, have you bought alcohol from a retailer (bar, restaurant, store, etc.) in the past **6 months**?

- A. Yes
- B. No

***** IF YOU ANSWERED "NO", SKIP TO QUESTION #18**

16. IF YOU ARE UNDER 21 and have bought alcohol from a retailer, which is the most common method? (Mark one.)

- A. Used a fake ID to buy alcohol
- B. Used someone else's ID to buy alcohol
- C. Bought alcohol without using an ID

17. IF YOU ARE UNDER 21 and have bought alcohol from a retailer, where are you most easily able to purchase alcohol? (Mark one.)

- A. Bar
- B. Restaurant
- C. Grocery store (e.g., Kroger, Marsh)
- D. Discount store (e.g., Wal-Mart, Target)
- E. Convenience store (e.g., Village Pantry, Circle K)
- F. Drug store (e.g., CVS)
- G. Liquor store
- H. Winery/microbrewery

18. Did you use other drugs (e.g., illegal drugs, misuse of medications) in the past **6 months**?

- A. Yes (go to question #19)
- B. No (go to question #20)

19. Have you experienced the following due to your use of drugs **in the past 6 months**. (Mark one for each line.)

	Number of Times					
	Never	Once	Twice	3-5	6-9	10 or more
a. Abused more than one drug at a time	A	B	C	D	E	F
b. Been unable to stop using drugs when you wanted to	A	B	C	D	E	F
c. Had blackouts or flashbacks as a result of drug use	A	B	C	D	E	F
d. Felt bad or guilty about your drug use	A	B	C	D	E	F
e. Had friends or family members worry or complain about your drug use	A	B	C	D	E	F
f. Created problems between you and your friends or family members	A	B	C	D	E	F
g. Engaged in illegal activities in order to obtain drugs	A	B	C	D	E	F
h. Gotten into physical fights because you were using drugs	A	B	C	D	E	F
i. Experienced withdrawal symptoms (felt sick) when you stopped taking drugs	A	B	C	D	E	F
j. Had medical problems as a result of your drug use (e.g. memory loss, infections, convulsions, bleeding)	A	B	C	D	E	F
k. Driven a car while under the influence of drugs	A	B	C	D	E	F
l. Gotten in trouble at work or school because of your drug use	A	B	C	D	E	F
m. Been in vulnerable situations (e.g., could have been hurt and/or date raped when using drugs)	A	B	C	D	E	F
n. Engaged in risky sexual behavior such as having unprotected sex or sex with someone you just met because you were using drugs	A	B	C	D	E	F
o. Failed a class because of drug use	A	B	C	D	E	F
p. Missed class or an assignment due to drug use	A	B	C	D	E	F

20. How do you think other students on campus feel (or would feel) about... (Mark one for each line.)

	Strongly approve	Somewhat approve	Neither approve or disapprove	Somewhat disapprove	Strongly disapprove
a. Trying marijuana once or twice	A	B	C	D	E
b. Smoking marijuana occasionally	A	B	C	D	E
c. Smoking marijuana regularly	A	B	C	D	E
d. Trying cocaine once or twice	A	B	C	D	E
e. Taking cocaine regularly	A	B	C	D	E
f. Trying amphetamines (e.g., Adderall, Ritalin) once or twice	A	B	C	D	E
g. Taking amphetamines regularly	A	B	C	D	E
h. Having one or two alcoholic drinks nearly every day	A	B	C	D	E
i. Having three or four alcoholic drinks nearly every day	A	B	C	D	E
j. Having five or more alcoholic drinks in one sitting	A	B	C	D	E
k. Taking steroids for body building or improved athletic performance	A	B	C	D	E

Use this definition for the next question.
One drink is a 12-ounce bottle of beer, a 5-ounce glass of wine, or
one shot of liquor straight or in a mixed drink.

21. How many alcoholic drinks do you think the typical student at your school had the last time he/she “partied/socialized”? (If you think the typical student at your school does not drink alcohol, enter 00. If less than 10, enter 01, 02, 03, etc.)

--	--

22. In the past year, how often have you done the following for *money*?

	Never	1-2 times/year	3-6 times/year	Monthly	Weekly	3 or more times/week
a. Played cards at home	A	B	C	D	E	F
b. Gambled at a casino	A	B	C	D	E	F
c. Gambled on-line	A	B	C	D	E	F
d. Gambled at informal locations (e.g., street corners, gambling houses)	A	B	C	D	E	F
e. Bet on fantasy football/basketball	A	B	C	D	E	F
f. Betting pools (e.g., March Madness)	A	B	C	D	E	F
g. Lottery	A	B	C	D	E	F
h. Placed bet with bookie	A	B	C	D	E	F

23. In the past year, if you gambled in ways other than online or playing the lottery, **where** did you most frequently gamble?

- A. I did not gamble in the past year
- B. I only gambled online or playing the lottery
- C. Apartment or house
- D. Residence hall
- E. Fraternity or sorority
- F. Other campus housing
- G. Casino
- H. Other

24. What consequences have you experienced due to your gambling?

	I do not gamble	None	Occasionally	Frequently
a. Academic problems (missed class, did poorly on exam, lower grades)	A	B	C	D
b. Issues with legal system	A	B	C	D
c. Financial issues	A	B	C	D
d. Felt bad about gambling	A	B	C	D
e. Poor physical health (sleep issues, depression, poor nutrition, poor hygiene)	A	B	C	D
f. Issues with friends and family (lied, strained relationships, arguments)	A	B	C	D

25. Thinking about your mental health, which includes stress, depression, and problems with emotions, for how many days during the past 30 days was your mental health not good?
(E.g., 00, 01, 02, 03, etc.)

--	--

__ Don't know / Not sure

26. On how many days in the past 2 weeks have you had thoughts that you would be better off dead or of hurting yourself in some way? (E.g., 00, 01, 02, 03, etc.)

--	--

27. Gender:

- M. Male
- F. Female

28. Do you consider yourself to be ...

- A. Heterosexual or straight
- B. Homosexual, gay, or lesbian
- C. Bisexual
- D. Other
- E. Don't Know/Not Sure/Questioning

29. Ethnicity:

- A. Hispanic
- B. Non-Hispanic

30. Race: (Mark one only.)

- A. White
- B. Black/African American
- C. Asian
- D. Native American/Alaskan Native
- E. Hawaiian/Pacific Islander
- F. More than one race
- G. Other

31. Marital status:

- A. Single, never married
- B. Single, living with partner
- C. Married
- D. Separated
- E. Divorced
- F. Widowed

32. Student status:

- A. Full-time
- B. Part-time

33. Classification:

- A. Freshman/First year
- B. Sophomore/Second year
- C. Junior
- D. Senior
- E. Grad/professional
- F. Not seeking a degree
- G. Other

34. Current residence as a student:

- A. On-campus
- B. Off-campus

35. Where do you live?

- A. House/apartment/etc.
- B. Residence hall
- C. Fraternity or sorority
- D. Other campus housing
- E. Other

36. With whom do you live? (Mark all that apply)

- A. Alone
- B. With roommate(s)
- C. With parent(s)
- D. With spouse
- E. With children
- F. Other

37. Are you employed?

- A. Yes, full-time
- B. Yes, part-time
- C. No

38. Have you ever served on active duty in the United States Armed Forces, either in the regular military or in a National Guard or military reserve unit? Active duty does not include training for the Reserves or National Guard, but DOES include activation, for example, for the Persian Gulf War.

- A. Yes
- B. No

Results of the Indiana College Substance Use Survey 2012

by
Rosemary King, M.P.H.
Mi Kyung Jun, Ph.D.
Junghun Lee, M.S.

*Funded by the Indiana Family and Social Services
Administration, Division of Mental Health and Addiction
through a contract with the Indiana Collegiate Action
Network*

Contact:
Indiana Prevention Resource Center
501 North Morton Street Suite 110
Bloomington, IN 47404
<http://www.drugs.indiana.edu>
drugprc@indiana.edu
Toll Free: 1-800-346-3077
Fax: 812-855-4940

